

**WO9602645**

Publication Title:

**HTK LIGAND**

Abstract:

A novel hepatoma transmembrane kinase receptor ligand (Htk ligand) which binds to, and activates, the Htk receptor is disclosed. As examples, mouse and human Htk ligands have been identified in a variety of tissues using a soluble Htk-Fc fusion protein. The ligands have been cloned and sequenced. The invention also relates to nucleic acids encoding the ligand, methods for production and use of the ligand, and antibodies directed thereto.

-----  
Data supplied from the esp@cenet database - <http://ep.espacenet.com>

*This Patent PDF Generated by Patent Fetcher(TM), a service of Stroke of Color, Inc.*

PCT

WORLD INTELLECTUAL  
PROPERTY ORGANIZATION

## INTERNATIONAL APPLICATION PUBLISHED

(51) International Patent Classification 6 :

C12N 15/12, C07K 14/47, A61K 38/17,  
C12N 15/62, C07K 19/00, 16/18, G01N  
33/68 // C07K 14/705

A2

WO 9602645A2

(43) International Publication Date: 1 February 1996 (01.02.96)

(21) International Application Number: PCT/US95/08812

(22) International Filing Date: 14 July 1995 (14.07.95)

(30) Priority Data:

08/277,722

20 July 1994 (20.07.94)

US

(71) Applicant: GENENTECH, INC. [US/US]; 460 Point San  
Bruno Boulevard, South San Francisco, CA 94080-4990  
(US).(72) Inventors: BENNETT, Brian, D.; 1332 Oddstad Boulevard,  
Pacifica, CA 94044 (US). MATTHEWS, William; 560  
Summit Springs Road, Woodside, CA 94062 (US).(74) Agents: LEE, Wendy, M. et al.; Genentech, Inc., 460 Point  
San Bruno Boulevard, South San Francisco, CA 94080-4990  
(US).(81) Designated States: AU, CA, JP, MX, European patent (AT,  
BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL,  
PT, SE).

Published

Without international search report and to be republished  
upon receipt of that report.

(54) Title: HTK LIGAND

```

      1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30
      31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60
      61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90
      91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120
      121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150
      151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180
      181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210
      211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240
      241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270
      271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300
      301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330
      331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360
      361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390
      391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420
      421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450
      451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480
      481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510
      511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540
      541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570
      571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600
      601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630
      631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660
      661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690
      691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720
      721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750
      751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780
      781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810
      811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840
      841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870
      871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900
      901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930
      931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960
      961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990
      991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020
      1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050
      1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080
      1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110
      1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140
      1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170
      1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200
      1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230
      1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260
      1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290
      1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320
      1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350
      1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380
      1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410
      1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440
      1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470
      1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500
      1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530
      1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560
      1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590
      1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620
      1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650
      1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680
      1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710
      1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740
      1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770
      1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800
      1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830
      1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860
      1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890
      1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920
      1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950
      1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980
      1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010
      2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040
      2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070
      2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100
      2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130
      2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160
      2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190
      2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220
      2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250
      2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280
      2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310
      2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340
      2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370
      2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400
      2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430
      2431 2432 
```

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

HTK LIGANDBACKGROUND OF THE INVENTIONField of the Invention

The present invention pertains generally to a receptor protein  
5 tyrosine kinase (rPTK) ligand. More particularly, the invention relates  
to a novel ligand which binds to, and activates, the hepatoma transmembrane  
kinase (Htk) receptor (also known as HpTK 5 receptor) and the isolation and  
recombinant production of the same.

Description of Related Art

10 Transduction of signals that regulate cell growth and differentiation  
is regulated in part by phosphorylation of various cellular proteins.  
Protein tyrosine kinases are enzymes that catalyze this process. Members  
of the protein tyrosine kinase family can be recognized by the presence of  
several conserved amino acid regions in the tyrosine kinase catalytic  
15 domain (Hanks et al., Science 241:42-52 [1988]). The tyrosine kinase  
domain is involved in the signal transduction pathways of mitogenesis,  
transformation and cell differentiation. Certain tyrosine kinases  
predominantly stimulate cell growth and differentiation, whereas other  
tyrosine kinases arrest growth and promote differentiation. Furthermore,  
20 depending on the cellular environment in which it is expressed, the same  
tyrosine kinase may either stimulate, or inhibit, cell proliferation. See  
Schlessinger et al., Neuron 9:383-391 [1992].

Receptor protein tyrosine kinases (rPTKs) convey extracellular  
signals to intracellular signaling pathways thereby controlling cell  
25 proliferation and differentiation. These rPTKs share a similar  
architecture, with an intracellular catalytic portion, a transmembrane  
domain and an extracellular ligand-binding domain. (Schlessinger et al.,  
*supra*). The extracellular domains (ECDs), which are responsible for ligand  
binding and transmission of biological signals, have been shown to be  
30 composed of a number of distinct structural motifs. The intracellular  
domain comprises a catalytic protein tyrosine kinase.

Receptor tyrosine kinases are categorized in several classes,  
according to sequence and structural similarities. For example, Class V  
receptors have cysteine rich and fibronectin Type III regions in the  
35 extracellular domain and include the EPH, ELK, ERK, BEK, ECK and HEK  
receptors. For a review of the various classes of receptor tyrosine  
kinases and their functions, see, e.g., Hanks et al., *supra* and  
Schlessinger et al., *supra*.

Protein ligands for receptor protein tyrosine kinases bind to the  
40 extracellular domain of their cognate receptors at the cell surface and  
thereby stimulate tyrosine phosphorylation. Several of these ligands are  
growth factors or cytokines, such as insulin-like growth factor 1 (IGF-1),  
epidermal growth factor (EGF), fibroblast growth factor (FGF), and nerve  
growth factor (NGF). Ligands for a number of tyrosine kinase receptors

have been shown to function within the hematopoietic system. For example, the ligand for murine flt3/flk-2 tyrosine kinase receptor, recently cloned, stimulates the proliferation of primitive mouse hematopoietic cells and human CD34-positive bone marrow cells. Lyman et al., Cell 75:1157-1167  
5 (1993).

A protein ligand which stimulates phosphorylation of the ECK receptor has recently been cloned and expressed in CHO cells. Bartley et al., Nature 368:558-560 (1994). This ECK ligand was found to be identical to B61, a molecule previously isolated by Holzman et al., Mol. Cell. Biol. 10:5830-  
10 5838 (1990).

A receptor tyrosine kinase has been recently identified and cloned from a human hepatocellular carcinoma cell line, Hep 3B. This receptor, called "Htk" receptor or "HPTK 5" receptor, is thought to belong to the Class V or EPH subfamily or rPTKs. See Bennett et al., J. Biol. Chem.,  
15 269(19):14211-14218 (1994).

Northern blot analysis of human fetal tissues revealed that expression of Htk receptor nucleic acid occurs in heart, lung, liver, brain and kidney. In adult human tissue, no signal was detectable in brain, while placenta had a particularly intense signal followed by kidney, liver,  
20 lung and pancreas. Skeletal muscle and heart were of lower signal intensity. See Bennett et al., *supra*.

Htk receptor nucleic acid expression in human tumor cell lines has also been analyzed by Northern blot analysis. Cell lines derived from liver, breast (MCF-7), colon (Colo 205), lung (NCI 69), melanocyte (HM-1)  
25 and cervix (HeLa) had detectable signals of appropriate size. Message was present in select cell lines of hematopoietic origin. K562 (a primitive myeloid cell with multipotential), THP-1 (a monocytoid cell), U937 (a myelomonocytic cell line), Hep3B (a human hepatocarcinoma cell line), and CMK (of megakaryocytic origin) were all positive for Htk receptor message,  
30 but lymphoid (H9, Jurkat, JH-1, Raji, Ramos) or select other myeloid cells (KG-1 or KMT2) had no detectable transcript by Northern analysis. See Bennett et al., *supra*.

The mouse homologue to the Htk receptor, called "myk-1", was isolated from mammary gland epithelia. See Andres et al., Oncogene 9:1461-1467  
35 (1994). Andres et al. report that myk-1 is induced during proliferation of mammary epithelium and down-regulated during its differentiation. Additionally, deregulated expression of the receptor is considered to potentially represent an early event in mammary gland carcinogenesis (see Andres et al., *supra*).

40 However, it is believed that the protein ligand for the Htk receptor has not been heretofore disclosed. Therefore, it is an object of the invention to provide a ligand to the Htk receptor.

It is a further object of the invention to provide nucleic acid encoding the Htk ligand so that the ligand can be made by recombinant DNA  
45 techniques.

These and other objects will be apparent to the ordinary artisan upon consideration of the specification as a whole.

#### SUMMARY OF THE INVENTION

These objects are accomplished, in one aspect, by providing isolated  
5 Htk ligand that may be antigenically or biologically active. In one embodiment, the invention provides a soluble form of the ligand with at least the transmembrane region deleted. Usually, the cytoplasmic domain will also be absent.

One example of a soluble form of the Htk ligand is an immunoadhesin  
10 which is a fusion of the extracellular domain of the Htk ligand and an immunoglobulin sequence.

The invention also pertains to other chimeras comprising the Htk ligand (or a portion thereof) fused to another polypeptide. An example of such a chimera is epitope tagged Htk ligand.

15 In another aspect, the invention provides a composition comprising biologically active Htk ligand and a pharmaceutically acceptable carrier. Preferably, the Htk ligand is present in a soluble form in the pharmaceutical composition.

The invention also provides isolated nucleic acid sequences encoding  
20 Htk ligand and Htk ligand chimeras.

The nucleic acid can be provided in a replicable vector which may be transformed into a host cell in one embodiment of the invention. A method of using the nucleic acid encoding the Htk ligand to effect the production of the novel protein is also provided which comprises expressing the  
25 nucleic acid in a culture of the transformed host cells and recovering the protein from the host cell culture.

The invention also provides a method which involves contacting the Htk receptor with the Htk ligand in order to cause phosphorylation of the kinase domain thereof.

30 The invention also provides a monoclonal antibody which binds to the Htk ligand, which can be used to detect the presence of the Htk ligand in a biological sample suspected of having the ligand, for example.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1B depict an alignment of the nucleotide sequence (SEQ ID  
35 NO: 1) and deduced amino acid sequence (SEQ ID NO: 2) of the murine Htk ligand described herein.

Figure 2 depicts an alignment of the nucleotide (SEQ ID NO: 3) and deduced amino acid sequence (SEQ ID NO: 4) of the human Htk ligand described herein.

40 Figure 3 shows an alignment of the amino acid sequences of murine Htk ligand (muHtkL) and human Htk ligand (humHtkL) (SEQ ID NOS: 2 and 4, respectively). Identical residues are enclosed within line boundaries. The shaded area represents a transmembrane domain. The extracellular domain and intracellular domain are N-terminal and C-terminal to the

transmembrane domain, respectively. The amino acid predicted to be the cleavage site for the signal peptide is indicated by an arrow. The potential N-linked glycosylation sites are marked with an (\*) and conserved cysteines are marked with an (v).

5 Figure 4 depicts the nucleotide sequence (SEQ ID NO: 5) and deduced amino acid sequence (SEQ ID NO: 6) of the human Htk receptor disclosed in Bennett et al., supra. The amino acid predicted to be the cleavage site for the signal peptide is indicated by an arrow. Cysteine residues conserved among ELK subfamily members are circled and the transmembrane  
10 region is overlined.

Figures 5A-5B show binding competition curves of Htk-Fc to the SV40MES 13 cell line (Figure 5A) or to recombinant murine Htk ligand expressed in COS-7 cells (Figure 5B). Scatchard representation of each binding curve is shown in the inset and revealed Kd's of 3 nM and 0.5 nM,  
15 respectively.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

##### I. Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

20 "Htk ligand" is defined herein to be any polypeptide sequence that binds to and activates a rPTK, preferably binds to the extracellular domain of the Htk receptor and thereby activates the intracellular tyrosine kinase domain thereof. Activation of the rPTK can be measured by autophosphorylation of tyrosine residues in the intracellular domain of the  
25 rPTK. See Example 4 herein for an exemplary technique for measuring receptor autophosphorylation. The Htk ligand may also may possess another biological property of a naturally occurring polypeptide, which polypeptide has either of the amino acid sequences shown in Figure 3.

"Biological property" for the purposes herein means an in vivo  
30 effector or antigenic function or activity that is directly or indirectly performed by the Htk ligand as shown by the sequences in Figure 3 (whether in its native or denatured conformation). A principal effector function is the ability of the Htk ligand to bind to, and activate, a rPTK such as the Htk receptor (also known as the Hptk 5 receptor), which is disclosed in  
35 Bennett et al., supra. The Htk receptor is a rPTK of the Class V or EPH subfamily of rPTKs. The nucleotide and amino acid sequence of the Htk receptor are depicted in Figure 4. Generally, the ligand will bind to the extracellular domain of the Htk receptor and thereby activate the intracellular tyrosine kinase domain thereof. Consequently, binding of the  
40 ligand to the receptor may result in enhancement or inhibition of proliferation and/or differentiation and/or activation of cells having a receptor for the Htk ligand in vivo or in vitro. Binding of the ligand to the Htk receptor can be determined using conventional techniques, including competitive binding methods, such as RIAs, ELISAs, and other competitive  
45 binding assays. Ligand/receptor complexes can be identified using such

separation methods as filtration, centrifugation, flow cytometry (see, e.g., Lyman et al., Cell 75:1157-1167 [1993]; Urdal et al., J. Biol. Chem. 263:2870-2877 [1988]; and Gearing et al., EMBO J 8:3667-3676 [1989]), and the like. Results from binding studies can be analyzed using any conventional graphical representation of the binding data, such as Scatchard analysis (Scatchard, Ann. NY Acad. Sci. 51:660-672 [1949]; and Goodwin et al., Cell 73:447-456 [1993]), and the like. Since the Htk ligand induces phosphorylation of the Htk receptor, conventional tyrosine phosphorylation assays, such as the assay described in Example 4 herein, can also be used as an indication of the formation of an Htk receptor/ligand complex. Other effector functions include signal transduction, any enzyme activity or enzyme modulatory activity (e.g., tyrosine kinase activity), or any structural role, for example. However, effector functions do not include possession of an epitope or antigenic site that is capable of cross-reacting with antibodies raised against Htk ligand. An antigenic function means possession of an epitope or antigenic site that is capable of cross-reacting with antibodies raised against the polypeptide sequence of a naturally occurring polypeptide comprising either of the polypeptide sequences of Figure 3.

"Biologically active" Htk ligand is defined herein as a polypeptide that shares an effector function of Htk ligand and that may (but need not) in addition possess an antigenic function. A principal known effector function of Htk ligand is its ability to cause protein phosphorylation of the Htk receptor.

"Antigenically active" Htk ligand is defined as a polypeptide that possesses an antigenic function of Htk ligand and that may (but need not) in addition possess an effector function.

In preferred embodiments, antigenically active Htk ligand is a polypeptide that binds with an affinity of at least about  $10^6$  l/mole to an antibody capable of binding Htk ligand. Ordinarily, the polypeptide binds with an affinity of at least about  $10^7$  l/mole. Isolated antibody capable of binding Htk ligand is an antibody that is identified and separated from a component of the natural environment in which it may be present. Most preferably, the antigenically active Htk ligand is a polypeptide that binds to an antibody capable of binding Htk ligand in its native conformation. Htk ligand in its native conformation is Htk ligand as found in nature that has not been denatured by chaotropic agents, heat, or other treatment that substantially modifies the three-dimensional structure of Htk ligand as determined, for example, by migration on non-reducing, non-denaturing sizing gels. Ordinarily, biologically or antigenically active Htk ligand will have an amino acid sequence having at least 75% amino acid sequence identity with either of the mature Htk ligand amino acid sequences shown in Figure 3, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, and most preferably at least 95%. Identity or homology with respect to this sequence is defined herein as the percentage of amino acid residues in the candidate sequence that are



identical with the Htk ligand residues, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. None of N-terminal, C-terminal, or internal extensions, deletions, or insertions into the Htk ligand sequence shall be construed as affecting sequence identity or homology.

Thus, the biologically active and antigenically active Htk ligand polypeptides that are the subject of this invention include the polypeptide represented by the entire translated nucleotide sequence of Htk ligand (including the signal sequence thereof); mature Htk ligand with the signal sequence cleaved therefrom; fragments consisting essentially of the intracellular domain or transmembrane domain of the Htk ligand; fragments of the Htk ligand having a consecutive sequence of at least 5, 10, 15, 20, 25, 30, or 40 amino acid residues from Htk ligand; amino acid sequence variants of Htk ligand wherein an amino acid residue has been inserted N- or C-terminal to, or within, Htk ligand or its fragment as defined above; amino acid sequence variants of Htk ligand or its fragment as defined above wherein an amino acid residue of Htk ligand or its fragment as defined above has been substituted by another residue, including predetermined mutations by, e.g., site-directed or PCR mutagenesis, Htk ligand of various animal species such as rabbit, rat, porcine, non-human primate, equine, murine, and ovine Htk ligand and alleles or other naturally occurring variants of the foregoing and human Htk ligand; derivatives of Htk ligand or its fragments as defined above wherein Htk ligand or its fragments have been covalent modified, by substitution, chemical, enzymatic, or other appropriate means, with a moiety other than a naturally occurring amino acid; and glycosylation variants of Htk ligand (insertion of a glycosylation site or alteration of any glycosylation site by deletion, insertion, or substitution of suitable residues). The preferred Htk ligand is human Htk ligand, especially native human Htk ligand having the sequence shown in Figure 2.

In one preferred embodiment, the Htk ligand comprises soluble Htk ligand. By "soluble Htk ligand" is meant Htk ligand which is essentially free of at least the transmembrane and, optionally, the intracellular domain of native Htk ligand. By "essentially free" is meant that the soluble Htk ligand sequence has less than 2% of the transmembrane domain, preferably 1.0-0% of the transmembrane domain, and more preferably 0.5-0% of this domain. The trans-membrane domains of the native murine and human amino acid sequences are delineated in Figure 3, i.e., residues 228 to 253 for murine Htk ligand and residues 225 to 250 for human Htk ligand. Such soluble Htk ligands may have advantages from a therapeutic standpoint because they are generally soluble in the patient's blood stream, for example. Similarly, such soluble ligands may prove to be particularly useful as diagnostics since they are expected to have a reduced tendency to incorporate in the cell membrane.

One example of a soluble form of the Htk ligand is an "immunoadhesin". The term "immunoadhesin" is used interchangeably with the expression "Htk ligand-immunoglobulin chimera" and refers to a chimeric molecule that combines the extracellular domain (ECD) of the Htk ligand with an immunoglobulin sequence. The immunoglobulin sequence preferably, but not necessarily, is an immunoglobulin constant domain. The immunoglobulin moiety in the chimeras of the present invention may be obtained from IgG-1, IgG-2, IgG-3 or IgG-4 subtypes, IgA, IgE, IgD or IgM, but preferably IgG-1 or IgG-3.

The expression "extracellular domain" or "ECD" when used herein refers to any polypeptide sequence that shares a receptor binding function of the extracellular domain of the naturally occurring Htk ligand disclosed herein. Receptor binding function refers to the ability of the polypeptide to bind the extracellular domain of a rPTK, such as the Htk receptor, and, optionally, activate the receptor. Accordingly, it is not necessary to include the entire extracellular domain since smaller segments are commonly found to be adequate for receptor binding. The term ECD encompasses polypeptide sequences in which the cytoplasmic domain and hydrophobic transmembrane sequence (and, optionally, 1-20 amino acids amino-terminal to the transmembrane domain) of the mature Htk ligand have been deleted. The extracellular domain of the Htk ligand is delineated in Figure 3 (i.e., it is amino-terminal to the transmembrane domain).

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising the entire Htk ligand, or a portion thereof, fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody thereagainst can be made, yet is short enough such that it does not interfere with activity of the Htk ligand. The tag polypeptide preferably also is fairly unique so that the antibody thereagainst does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least 6 amino acid residues and usually between about 8-50 amino acid residues (preferably between about 9-30 residues).

An "exogenous" therapeutic compound is defined herein to mean a therapeutic compound that is foreign to the mammalian patient, or homologous to a compound found in the mammalian patient but produced outside the mammalian patient.

"Isolated", when used to describe the various proteins disclosed herein, means protein that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the protein, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the protein will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue

or, preferably, silver stain. Isolated protein includes protein in situ within recombinant cells, since at least one component of the Htk ligand natural environment will not be present. Ordinarily, however, isolated protein will be prepared by at least one purification step.

5 "Essentially pure" protein means a composition comprising at least about 90% by weight of the protein, based on total weight of the composition, preferably at least about 95% by weight. "Essentially homogeneous" protein means a composition comprising at least about 99% by weight of protein, based on total weight of the composition.

10 In accordance with this invention, "Htk ligand nucleic acid" or a "Htk ligand nucleic acid molecule" is RNA or DNA containing greater than ten bases that encodes a biologically active or antigenically active Htk ligand, is complementary to nucleic acid sequence encoding such Htk ligand, or hybridizes to nucleic acid sequence encoding such Htk ligand and remains  
15 stably bound to it under stringent conditions. The nucleic acid optionally includes the regions of the nucleic acid sequences of Figure 1A and Figure 2 which encode the signal sequences. In one embodiment, the nucleic acid sequence is selected from:

- 20 (a) the coding regions of the nucleic acid sequences of Figure 1A or Figure 2;
- (b) a sequence corresponding to either of the sequences of (a) within the scope of degeneracy of the genetic code; or
- (c) a sequence which hybridizes with a sequence complementary to the sequences of (a) or (b) under stringent conditions and  
25 which codes for a biologically active Htk ligand.

In one preferred embodiment, the nucleic acid encodes soluble Htk ligand wherein the nucleic acid encoding the transmembrane region, and optionally the cytoplasmic region, of the polypeptide has been deleted.

30 Preferably, the Htk ligand nucleic acid molecule encodes a polypeptide sharing at least 75% sequence identity, more preferably at least 80%, still more preferably at least 85%, even more preferably at least 90%, and most preferably 95%, with either of the Htk ligand amino acid sequences shown in Figure 3. Preferably, the Htk ligand nucleic acid molecule that hybridizes to nucleic acid sequence encoding Htk ligand  
35 contains at least 20, more preferably 40, and most preferably 90 bases.

"Stringent conditions" are those that (1) employ low ionic strength and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate/0.1% NaDodSO<sub>4</sub> at 50°C; (2) employ during hybridization a denaturing agent such as formamide, for example, 50% (vol/vol) formamide with 0.1%  
40 bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran  
45 sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS.

An "isolated" Htk ligand nucleic acid molecule is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the Htk ligand nucleic acid. An isolated Htk ligand nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated Htk ligand nucleic acid molecules therefore are distinguished from the Htk ligand nucleic acid molecule as it exists in natural cells. However, an isolated Htk ligand nucleic acid molecule includes Htk ligand nucleic acid molecules contained in cells that ordinarily express Htk ligand where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The isolated Htk ligand polypeptide, Htk ligand nucleic acid, or Htk ligand antibody may be labeled for diagnostic and probe purposes, using a label as described and defined further below in the discussion on uses of Htk ligand antibodies.

The expression "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, a ribosome binding site, and possibly, other as yet poorly understood sequences. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accord with conventional practice.

The term "antibody" is used in the broadest sense and specifically covers single anti-Htk ligand monoclonal antibodies (including agonist and antagonist antibodies) and anti-Htk ligand antibody compositions with polypepitopic specificity.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional

(polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen.

5       The monoclonal antibodies herein include hybrid and recombinant antibodies produced by splicing a variable (including hypervariable) domain of an anti-Htk ligand antibody with a constant domain (e.g. "humanized" antibodies), or a light chain with a heavy chain, or a chain from one species with a chain from another species, or fusions with heterologous  
10 proteins, regardless of species of origin or immunoglobulin class or subclass designation, as well as antibody fragments (e.g., Fab, F(ab')<sub>2</sub>, and Fv), so long as they exhibit the desired biological activity. [See, e.g. US Pat No 4,816,567 and Mage & Lamoyi, in Monoclonal Antibody Production Techniques and Applications, pp.79-97 (Marcel Dekker, Inc., New York  
15 (1987))].

      Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies  
20 to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler & Milstein, Nature 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage libraries generated using the techniques described in McCafferty et al.,  
25 Nature 348:552-554 (1990), for example.

      "Humanized" forms of non-human (e.g. murine) antibodies are specific chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub>, or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human  
30 immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances,  
35 Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In general, the  
40 humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise  
45 at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

## II. Modes for Practicing Out the Invention

The present invention is based on the discovery of a novel Htk ligand which binds to, and activates, the Htk receptor.

The murine Htk ligand cDNA sequence is depicted in Figures

5 1A-B. The predicted molecular weight of the protein following signal peptide cleavage is 34 kD with an estimated pI of 8.9. Similarly, the human Htk ligand has been identified and isolated. The nucleotide and amino acid sequences of the human Htk ligand are shown in Figure 2. The murine and human ligands show 96% homology at the amino acid level,  
10 demonstrating a high degree of conservation between species. A description follows as to the preparation of Htk ligand and variants thereof.

### 1. Preparation of Natural Sequence Htk Ligand and Variants Thereof

Most of the discussion below pertains to production of Htk ligand by culturing cells transformed with a vector containing Htk ligand nucleic  
15 acid and recovering the polypeptide from the cell culture. It is further envisioned that the Htk ligand of this invention may be produced by homologous recombination, as provided for in WO 91/06667, published 16 May 1991.

Briefly, this method involves transforming primary mammalian cells  
20 containing endogenous Htk ligand gene (e.g., human cells if the desired Htk ligand is human) with a construct (i.e., vector) comprising an amplifiable gene [such as dihydrofolate reductase (DHFR) or others discussed below] and at least one flanking region of a length of at least about 150 bp that is homologous with a DNA sequence at the locus of the coding region of the Htk  
25 ligand gene to provide amplification of the Htk ligand gene. The amplifiable gene must be at a site that does not interfere with expression of the Htk ligand gene. The transformation is conducted such that the construct becomes homologously integrated into the genome of the primary cells to define an amplifiable region.

30 Primary cells comprising the construct are then selected for by means of the amplifiable gene or other marker present in the construct. The presence of the marker gene establishes the presence and integration of the construct into the host genome. No further selection of the primary cells need be made, since selection will be made in the second host. If desired,  
35 the occurrence of the homologous recombination event can be determined by employing PCR and either sequencing the resulting amplified DNA sequences or determining the appropriate length of the PCR fragment when DNA from correct homologous integrants is present and expanding only those cells containing such fragments. Also if desired, the selected cells may be  
40 amplified at this point by stressing the cells with the appropriate amplifying agent (such as methotrexate if the amplifiable gene is DHFR), so that multiple copies of the target gene are obtained. Preferably, however, the amplification step is not conducted until after the second transformation described below.

After the selection step, DNA portions of the genome, sufficiently large to include the entire amplifiable region, are isolated from the selected primary cells. Secondary mammalian expression host cells are then transformed with these genomic DNA portions and cloned, and clones are selected that contain the amplifiable region. The amplifiable region is then amplified by means of an amplifying agent if not already amplified in the primary cells. Finally, the secondary expression host cells now comprising multiple copies of the amplifiable region containing Htk ligand are grown so as to express the gene and produce the protein.

10       A.     Isolation of DNA Encoding Htk Ligand

The DNA encoding Htk ligand may be obtained from any cDNA library prepared from tissue believed to possess the Htk ligand mRNA and to express it at a detectable level. Accordingly, human Htk ligand DNA can be conveniently obtained from a cDNA library prepared from human fetal lung or brain tissue. The murine Htk ligand DNA can be derived from a cDNA library of the SV40MES 13 cell line, for example. The Htk ligand gene may also be obtained from a genomic library or by oligonucleotide synthesis.

Libraries are screened with probes (such as antibodies to the Htk ligand or oligonucleotides of about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures as described in chapters 10-12 of Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding Htk ligand is to use PCR methodology as described in section 14 of Sambrook et al., *supra*.

A preferred method of practicing this invention is to use carefully selected oligonucleotide sequences to screen cDNA libraries from various tissues, preferably mammalian fetal lung or brain lines, more preferably, human fetal lung or brain cell lines. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized.

The oligonucleotide must be labeled such that it can be detected upon hybridization to DNA in the library being screened. The preferred method of labeling is to use <sup>32</sup>P- labeled ATP with polynucleotide kinase, as is well known in the art, to radiolabel the oligonucleotide. However, other methods may be used to label the oligonucleotide, including, but not limited to, biotinylation or enzyme labeling.

Of particular interest is the Htk ligand nucleic acid that encodes a full-length polypeptide. In some preferred embodiments, the nucleic acid sequence includes the native Htk ligand signal sequence. Nucleic acid having all the protein coding sequence is obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in section 7.79 of Sambrook et al.,

supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

B. Amino Acid Sequence Variants of Native Htk Ligand

Amino acid sequence variants of Htk ligand are prepared by  
5 introducing appropriate nucleotide changes into the Htk ligand DNA, or by  
synthesis of the desired Htk ligand polypeptide. Such variants include,  
for example, deletions from, or insertions or substitutions of, residues  
within the amino acid sequences shown for the Htk ligands in Figure 3. Any  
combination of deletion, insertion, and substitution is made to arrive at  
10 the final construct, provided that the final construct possesses the  
desired characteristics. The amino acid changes also may alter post-  
translational processes of the Htk ligand, such as changing the number or  
position of glycosylation sites, altering the membrane anchoring  
characteristics, and/or altering the intracellular location of the Htk  
15 ligand by inserting, deleting, or otherwise affecting the leader sequence  
of the Htk ligand.

For the design of amino acid sequence variants of Htk ligand, the  
location of the mutation site and the nature of the mutation will depend  
on the Htk ligand characteristic(s) to be modified. The sites for mutation  
20 can be modified individually or in series, e.g., by (1) substituting first  
with conservative amino acid choices and then with more radical selections  
depending upon the results achieved, (2) deleting the target residue, or  
(3) inserting residues of the same or a different class adjacent to the  
located site, or combinations of options 1-3.

25 A useful method for identification of certain residues or regions of  
the Htk ligand polypeptide that are preferred locations for mutagenesis is  
called "alanine scanning mutagenesis," as described by Cunningham and  
Wells, Science, 244:1081-1085 (1989). Here, a residue or group of target  
residues are identified (e.g., charged residues such as arg, asp, his, lys,  
30 and glu) and replaced by a neutral or negatively charged amino acid (most  
preferably alanine or polyalanine) to affect the interaction of the amino  
acids with the surrounding aqueous environment in or outside the cell.  
Those domains demonstrating functional sensitivity to the substitutions  
then are refined by introducing further or other variants at or for the  
35 sites of substitution. Thus, while the site for introducing an amino acid  
sequence variation is predetermined, the nature of the mutation per se need  
not be predetermined. For example, to optimize the performance of a  
mutation at a given site, ala scanning or random mutagenesis is conducted  
at the target codon or region and the expressed Htk ligand variants are  
40 screened for the optimal combination of desired activity.

There are two principal variables in the construction of amino acid  
sequence variants: the location of the mutation site and the nature of the  
mutation. These are variants of the sequences of Figure 3, and may  
represent naturally occurring alleles (which will not require manipulation  
45 of the Htk ligand DNA) or predetermined mutant forms made by mutating the  
DNA, either to arrive at an allele or a variant not found in nature. In



general, the location and nature of the mutation chosen will depend upon the Htk ligand characteristic to be modified. Obviously, such variations that, for example, convert Htk ligand into a known receptor protein tyrosine kinase ligand are not included within the scope of this invention.

5 Amino acid sequence deletions generally range from about 1 to 30 residues, more preferably about 1 to 10 residues, and typically are contiguous. Contiguous deletions ordinarily are made in even numbers of residues, but single or odd numbers of deletions are within the scope hereof. Deletions may be introduced into regions of low homology among Htk  
10 ligand and known Htk ligands (which share the most sequence identity to the human Htk ligand amino acid sequence) to modify the activity of Htk ligand. Deletions from Htk ligand in areas of substantial homology with homologous Htk ligand proteins will be more likely to modify the biological activity of Htk ligand more significantly. The number of consecutive deletions will  
15 be selected so as to preserve the tertiary structure of Htk ligand in the affected domain, e.g., beta-pleated sheet or alpha helix.

One preferred deletional variant is the soluble Htk ligand defined herein. This variant of the Htk ligand has the transmembrane and, optionally, intracellular domains deleted using the techniques for  
20 generating deletional variants.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions (i.e.,  
25 insertions within the mature Htk ligand sequence) may range generally from about 1 to 10 residues, more preferably 1 to 5, most preferably 1 to 3. Insertions are preferably made in even numbers of residues, but this is not required. Examples of terminal insertions include mature Htk ligand with an N-terminal methionyl residue, an artifact of the direct expression of  
30 mature Htk ligand in recombinant cell culture, and fusion of a heterologous N-terminal signal sequence to the N-terminus of the mature Htk ligand molecule to facilitate the secretion of mature Htk ligand from recombinant hosts. Such signal sequences generally will be obtained from, and thus homologous to, the intended host cell species. Suitable sequences include  
35 STII or lpp for *E. coli*, alpha factor or invertase for yeast, and viral signals such as herpes gD for mammalian cells.

Other insertional variants of the Htk ligand molecule include the fusion to the N- or C-terminus of Htk ligand of immunogenic polypeptides, e.g., bacterial polypeptides such as beta-lactamase or an enzyme encoded  
40 by the *E. coli* trp locus, or yeast protein, and C-terminal fusions with proteins having a long half-life such as immunoglobulin constant regions (or other immunoglobulin regions), albumin, or ferritin, as described in WO 89/02922 published 6 April 1989.

A third group of variants are amino acid substitution variants.  
45 These variants have at least one amino acid residue in the Htk ligand molecule removed and a different residue inserted in its place. The sites

of greatest interest for substitutional mutagenesis include sites identified as the active site(s) of Htk ligand and sites where the amino acids found in the known analogues are substantially different in terms of side-chain bulk, charge, or hydrophobicity, but where there is also a high degree of sequence identity at the selected site within various animal Htk ligand species. Other sites of interest are those in which particular residues of the Htk ligand obtained from various species are identical. These sites, especially those falling within a sequence of at least three other identically conserved sites, are substituted in a relatively conservative manner. Such conservative substitutions are shown in Table 1 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 1, or as further described below in reference to amino acid classes, are introduced and the products screened.

TABLE 1

	<u>Original Residue</u>	<u>Exemplary Substitutions</u>	<u>Preferred Substitutions</u>
	Ala (A)	val; leu; ile	val
20	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
	Cys (C)	ser	ser
	Gln (Q)	asn	asn
25	Glu (E)	asp	asp
	Gly (G)	pro	pro
	His (H)	asn; gln; lys; arg	arg
	Ile (I)	leu; val; met; ala; phe; norleucine	leu
30	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
	Met (M)	leu; phe; ile	leu
	Phe (F)	leu; val; ile; ala	leu
35	Pro (P)	gly	gly
	Ser (S)	thr	thr
	Thr (T)	ser	ser
	Trp (W)	tyr	tyr
	Tyr (Y)	trp; phe; thr; ser	phe
40	Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the Htk ligand are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a

sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- 5 (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- 10 (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

- 15 In one embodiment of the invention, it is desirable to inactivate one or more internal protease cleavage sites that are present in the molecule. These sites are identified by inspection of the encoded amino acid sequence, in the case of trypsin, e.g., for an arginyl or lysinyl residue. When protease cleavage sites are identified, they are rendered inactive to
- 20 proteolytic cleavage by substituting the targeted residue with another residue, preferably a basic residue such as glutamine or a hydrophobic residue such as serine; by deleting the residue; or by inserting a prolyl residue immediately after the residue.

- In another embodiment, any methionyl residues other than the starting
- 25 methionyl residue of the signal sequence, or any residue located within about three residues N- or C-terminal to each such methionyl residue, is substituted by another residue (preferably in accord with Table 1) or deleted. Alternatively, about 1-3 residues are inserted adjacent to such sites.

- 30 Any cysteine residues not involved in maintaining the proper conformation of Htk ligand also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking.

- Nucleic acid molecules encoding amino acid sequence variants of Htk
- 35 ligand are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a
- 40 non-variant version of Htk ligand.

- Oligonucleotide-mediated mutagenesis is a preferred method for preparing substitution, deletion, and insertion variants of Htk ligand DNA. This technique is well known in the art as described by Adelman et al., DNA, 2:183 (1983). Briefly, Htk ligand DNA is altered by hybridizing an
- 45 oligonucleotide encoding the desired mutation to a DNA template, where the template is the single-stranded form of a plasmid or bacteriophage

containing the unaltered or native DNA sequence of Htk ligand. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the Htk ligand DNA.

Generally, oligonucleotides of at least 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using techniques known in the art such as that described by Crea et al., Proc. Natl. Acad. Sci. USA 75:5765 (1978).

The DNA template can be generated by those vectors that are either derived from bacteriophage M13 vectors (the commercially available M13mp18 and M13mp19 vectors are suitable), or those vectors that contain a single-stranded phage origin of replication as described by Viera et al., Meth. Enzymol. 153:3 (1987). Thus, the DNA that is to be mutated may be inserted into one of these vectors to generate single-stranded template. Production of the single-stranded template is described in Sections 4.21-4.41 of Sambrook et al., *supra*.

Alternatively, single-stranded DNA template may be generated by denaturing double-stranded plasmid (or other) DNA using standard techniques.

For alteration of the native DNA sequence (to generate amino acid sequence variants, for example), the oligonucleotide is hybridized to the single-stranded template under suitable hybridization conditions. A DNA polymerizing enzyme, usually the Klenow fragment of DNA polymerase I, is then added to synthesize the complementary strand of the template using the oligonucleotide as a primer for synthesis. A heteroduplex molecule is thus formed such that one strand of DNA encodes the mutated form of Htk ligand, and the other strand (the original template) encodes the native, unaltered sequence of Htk ligand. This heteroduplex molecule is then transformed into a suitable host cell, usually a prokaryote such as *E. coli* JM101. After the cells are grown, they are plated onto agarose plates and screened using the oligonucleotide primer radiolabeled with <sup>32</sup>P to identify the bacterial colonies that contain the mutated DNA. The mutated region is then removed and placed in an appropriate vector for protein production, generally an expression vector of the type typically employed for transformation of an appropriate host.

The method described immediately above may be modified such that a homoduplex molecule is created wherein both strands of the plasmid contain the mutation(s). The modifications are as follows: The single-stranded oligonucleotide is annealed to the single-stranded template as described above. A mixture of three deoxyribonucleotides, deoxyriboadenosine (dATP), deoxyriboguanosine (dGTP), and deoxyribothymidine (dTTP), is combined with

a modified thio-deoxyribocytosine called dCTP-(aS) (which can be obtained from the Amersham Corporation). This mixture is added to the template-oligonucleotide complex. Upon addition of DNA polymerase to this mixture, a strand of DNA identical to the template except for the mutated  
5 bases is generated. In addition, this new strand of DNA will contain dCTP-(aS) instead of dCTP, which serves to protect it from restriction endonuclease digestion.

After the template strand of the double-stranded heteroduplex is nicked with an appropriate restriction enzyme, the template strand can be  
10 digested with ExoIII nuclease or another appropriate nuclease past the region that contains the site(s) to be mutagenized. The reaction is then stopped to leave a molecule that is only partially single-stranded. A complete double-stranded DNA homoduplex is then formed using DNA polymerase in the presence of all four deoxyribonucleotidetriphosphates, ATP, and DNA  
15 ligase. This homoduplex molecule can then be transformed into a suitable host cell such as *E. coli* JM101, as described above.

DNA encoding Htk ligand mutants with more than one amino acid to be substituted may be generated in one of several ways. If the amino acids are located close together in the polypeptide chain, they may be mutated  
20 simultaneously using one oligonucleotide that codes for all of the desired amino acid substitutions. If, however, the amino acids are located some distance from each other (separated by more than about ten amino acids), it is more difficult to generate a single oligonucleotide that encodes all of the desired changes. Instead, one of two alternative methods may be  
25 employed.

In the first method, a separate oligonucleotide is generated for each amino acid to be substituted. The oligonucleotides are then annealed to the single-stranded template DNA simultaneously, and the second strand of DNA that is synthesized from the template will encode all of the desired  
30 amino acid substitutions.

The alternative method involves two or more rounds of mutagenesis to produce the desired mutant. The first round is as described for the single mutants: wild-type DNA is used for the template, an oligonucleotide encoding the first desired amino acid substitution(s) is annealed to this  
35 template, and the heteroduplex DNA molecule is then generated. The second round of mutagenesis utilizes the mutated DNA produced in the first round of mutagenesis as the template. Thus, this template already contains one or more mutations. The oligonucleotide encoding the additional desired amino acid substitution(s) is then annealed to this template, and the  
40 resulting strand of DNA now encodes mutations from both the first and second rounds of mutagenesis. This resultant DNA can be used as a template in a third round of mutagenesis, and so on.

PCR mutagenesis is also suitable for making amino acid variants of Htk ligand. While the following discussion refers to DNA, it is understood  
45 that the technique also finds application with RNA. The PCR technique generally refers to the following procedure (see Erlich, Science, 252:1643-

1650 (1991), the chapter by R. Higuchi, p. 61-70). When small amounts of template DNA are used as starting material in a PCR, primers that differ slightly in sequence from the corresponding region in a template DNA can be used to generate relatively large quantities of a specific DNA fragment that differs from the template sequence only at the positions where the primers differ from the template. For introduction of a mutation into a plasmid DNA, one of the primers is designed to overlap the position of the mutation and to contain the mutation; the sequence of the other primer must be identical to a stretch of sequence of the opposite strand of the plasmid, but this sequence can be located anywhere along the plasmid DNA. It is preferred, however, that the sequence of the second primer is located within 200 nucleotides from that of the first, such that in the end the entire amplified region of DNA bounded by the primers can be easily sequenced. PCR amplification using a primer pair like the one just described results in a population of DNA fragments that differ at the position of the mutation specified by the primer, and possibly at other positions, as template copying is somewhat error-prone.

If the ratio of template to product material is extremely low, the vast majority of product DNA fragments incorporate the desired mutation(s). This product material is used to replace the corresponding region in the plasmid that served as PCR template using standard DNA technology. Mutations at separate positions can be introduced simultaneously by either using a mutant second primer, or performing a second PCR with different mutant primers and ligating the two resulting PCR fragments simultaneously to the vector fragment in a three (or more)-part ligation.

In a specific example of PCR mutagenesis, template plasmid DNA (1  $\mu$ g) is linearized by digestion with a restriction endonuclease that has a unique recognition site in the plasmid DNA outside of the region to be amplified. Of this material, 100 ng is added to a PCR mixture containing PCR buffer, which contains the four deoxynucleotide triphosphates and is included in the GeneAmp® kits (obtained from Perkin-Elmer Cetus, Norwalk, CT and Emeryville, CA), and 25 pmole of each oligonucleotide primer, to a final volume of 50  $\mu$ l. The reaction mixture is overlaid with 35  $\mu$ l mineral oil. The reaction mixture is denatured for five minutes at 100°C, placed briefly on ice, and then 1  $\mu$ l *Thermus aquaticus* (Taq) DNA polymerase (5 units/ $\mu$ l, purchased from Perkin-Elmer Cetus) is added below the mineral oil layer. The reaction mixture is then inserted into a DNA Thermal Cycler (purchased from Perkin-Elmer Cetus) programmed as follows:

2 min. 55°C  
30 sec. 72°C, then 19 cycles of the following:  
30 sec. 94°C  
30 sec. 55°C, and  
30 sec. 72°C.

At the end of the program, the reaction vial is removed from the thermal cycler and the aqueous phase transferred to a new vial, extracted with phenol/chloroform (50:50 vol), and ethanol precipitated, and the DNA

is recovered by standard procedures. This material is subsequently subjected to the appropriate treatments for insertion into a vector.

Another method for preparing variants, cassette mutagenesis, is based on the technique described by Wells et al., Gene, 34:315 (1985). The starting material is the plasmid (or other vector) comprising the Htk ligand DNA to be mutated. The codon(s) in the Htk ligand DNA to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate locations in the Htk ligand DNA. After the restriction sites have been introduced into the plasmid, the plasmid is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using standard procedures. The two strands are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 3' and 5' ends that are compatible with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated Htk ligand DNA sequence.

#### C. Insertion of Nucleic Acid into Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding native or variant Htk ligand is inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. Many vectors are available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

##### (i) Signal Sequence Component

The Htk ligands of this invention may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which is preferably a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the Htk ligand DNA that is inserted into the vector. The heterologous signal sequence selected preferably is one that is recognized and processed (i.e., cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the native Htk ligand signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the native signal sequence may be substituted by, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces*  $\alpha$ -factor leaders, the latter described in U.S. Pat. No. 5,010,182 issued 23 April 1991), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179

published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression the native signal sequence (e.g., the Htk ligand presequence that normally directs secretion of Htk ligand from human cells *in vivo*) is satisfactory, although other mammalian  
5 signal sequences may be suitable, such as signal sequences from other animal Htk ligands, and signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders, for example, the herpes simplex gD signal.

The DNA for such precursor region is ligated in reading frame to DNA  
10 encoding the mature Htk ligand.

(ii) Origin of Replication Component

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector  
15 to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 $\mu$  plasmid origin is suitable for yeast, and  
20 various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

25 Most expression vectors are "shuttle" vectors, i.e., they are capable of replication in at least one class of organisms but can be transfected into another organism for expression. For example, a vector is cloned in *E. coli* and then the same vector is transfected into yeast or mammalian cells for expression even though it is not capable of replicating  
30 independently of the host cell chromosome.

DNA may also be amplified by insertion into the host genome. This is readily accomplished using *Bacillus* species as hosts, for example, by including in the vector a DNA sequence that is complementary to a sequence found in *Bacillus* genomic DNA. Transfection of *Bacillus* with this vector  
35 results in homologous recombination with the genome and insertion of Htk ligand DNA. However, the recovery of genomic DNA encoding Htk ligand is more complex than that of an exogenously replicated vector because restriction enzyme digestion is required to excise the Htk ligand DNA.

(iii) Selection Gene Component

40 Expression and cloning vectors should contain a selection gene, also termed a selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection  
45 genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b)



complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

One example of a selection scheme utilizes a drug to arrest growth  
5 of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin (Southern et al., J. Molec. Appl. Genet. 1:327 [1982]), mycophenolic acid (Mulligan et al., Science 209:1422 [1980]) or hygromycin  
10 (Sugden et al., Mol. Cell. Biol. 5:410-413 [1985]). The three examples given above employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid), or hygromycin, respectively.

Another example of suitable selectable markers for mammalian cells  
15 are those that enable the identification of cells competent to take up the Htk ligand nucleic acid, such as DHFR or thymidine kinase. The mammalian cell transformants are placed under selection pressure that only the transformants are uniquely adapted to survive by virtue of having taken up the marker. Selection pressure is imposed by culturing the transformants  
20 under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes Htk ligand. Amplification is the process by which genes in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes  
25 of successive generations of recombinant cells. Increased quantities of Htk ligand are synthesized from the amplified DNA. Other examples of amplifiable genes include metallothionein-I and -II, preferably primate metallothionein genes, adenosine deaminase, ornithine decarboxylase, etc.

For example, cells transformed with the DHFR selection gene are first  
30 identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity, prepared and propagated as described by Urlaub and Chasin, Proc. Natl. Acad. Sci. USA  
35 77:4216 (1980). The transformed cells are then exposed to increased levels of methotrexate. This leads to the synthesis of multiple copies of the DHFR gene, and, concomitantly, multiple copies of other DNA comprising the expression vectors, such as the DNA encoding Htk ligand. This amplification technique can be used with any otherwise suitable host, e.g.,  
40 ATCC No. CCL61 CHO-K1, notwithstanding the presence of endogenous DHFR if, for example, a mutant DHFR gene that is highly resistant to Mtx is employed (EP 117,060).

Alternatively, host cells [particularly wild-type hosts that contain endogenous DHFR] transformed or co-transformed with DNA sequences encoding  
45 Htk ligand, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth

in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, e.g., kanamycin, neomycin, or G418. See U.S. Patent No. 4,965,199.

A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 (Stinchcomb et al., Nature 282:39 [1979]; Kingsman et al., Gene 7:141 [1979]; or Tschemper et al., Gene 10:157 [1980]). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 (Jones, Genetics 85:12 [1977]). The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, *Leu2*-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu2* gene.

In addition, vectors derived from the 1.6  $\mu$ m circular plasmid pKD1 can be used for transformation of *Kluyveromyces* yeasts. Bianchi et al., Curr. Genet. 12:185 (1987). More recently, an expression system for large-scale production of recombinant calf chymosin was reported for *K. lactis*. Van den Berg, Bio/Technology 8:135 (1990). Stable multi-copy expression vectors for secretion of mature recombinant human serum albumin by industrial strains of *Kluyveromyces* have also been disclosed. Fleer et al., Bio/Technology 9:968-975 (1991).

(iv) Promoter Component

Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the Htk ligand nucleic acid. Promoters are untranslated sequences located upstream (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of particular nucleic acid sequence, such as the Htk ligand nucleic acid sequence, to which they are operably linked. Such promoters typically fall into two classes, inducible and constitutive. Inducible promoters are promoters that initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, e.g., the presence or absence of a nutrient or a change in temperature. At this time a large number of promoters recognized by a variety of potential host cells are well known. These promoters are operably linked to Htk ligand-encoding DNA by removing the promoter from the source DNA by restriction enzyme digestion and inserting the isolated promoter sequence into the vector. Both the native Htk ligand promoter sequence and many heterologous promoters may be used to direct amplification and/or expression of the Htk ligand DNA. However, heterologous promoters are preferred, as they generally permit greater transcription and higher yields of Htk ligand as compared to the native Htk ligand promoter.

Promoters suitable for use with prokaryotic hosts include the  $\beta$ -lactamase and lactose promoter systems (Chang et al., Nature 275:615 [1978]; and Goeddel et al., Nature 281:544 [1979]), alkaline phosphatase, a tryptophan (*trp*) promoter system (Goeddel, Nucleic Acids Res., 8:4057

[1980] and EP 36,776) and hybrid promoters such as the tac promoter (deBoer et al., Proc. Natl. Acad. Sci. USA 80:21-25 [1983]). However, other known bacterial promoters are suitable. Their nucleotide sequences have been published, thereby enabling a skilled worker operably to ligate them to DNA encoding Htk ligand (Siebenlist et al., Cell 20:269 [1980]) using linkers or adaptors to supply any required restriction sites. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding Htk ligand.

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CXCAAT region where X may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman et al., J. Biol. Chem. 255:2073 [1980]) or other glycolytic enzymes (Hess et al., J. Adv. Enzyme Reg. 7:149 [1968]; and Holland, Biochemistry 17:4900 [1978]), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in Hitzeman et al., EP 73,657A. Yeast enhancers also are advantageously used with yeast promoters.

Htk ligand transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, from heat-shock promoters, and from the promoter normally associated with the Htk ligand sequence, provided such promoters are compatible with the host cell systems.

The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. Fiers et al., Nature 273:113 (1978); Mulligan and Berg, Science 209:1422-1427 (1980); Pavlakis et al., Proc. Natl. Acad. Sci.

USA 78:7398-7402 (1981). The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. Greenaway et al., Gene 18:355-360 (1982). A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Patent No. 4,419,446. A modification of this system is described in U.S. Patent No. 4,601,978. See also Gray et al., Nature 295:503-508 (1982) on expressing cDNA encoding immune interferon in monkey cells; Reyes et al., Nature 297:598-601 (1982) on expression of human  $\beta$ -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus; Canaani and Berg, Proc. Natl. Acad. Sci. USA 79:5166-5170 (1982) on expression of the human interferon  $\beta$ 1 gene in cultured mouse and rabbit cells; and Gorman et al., Proc. Natl. Acad. Sci. USA 79:6777-6781 (1982) on expression of bacterial CAT sequences in CV-1 monkey kidney cells, chicken embryo fibroblasts, Chinese hamster ovary cells, HeLa cells, and mouse NIH-3T3 cells using the Rous sarcoma virus long terminal repeat as a promoter.

(v) Enhancer Element Component

Transcription of a DNA encoding the Htk ligand of this invention by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Enhancers are relatively orientation and position independent, having been found 5' (Laimins et al., Proc. Natl. Acad. Sci. USA 78:993 [1981]) and 3' (Lusky et al., Mol. Cell Bio. 3:1108 [1983]) to the transcription unit, within an intron (Banerji et al., Cell 33:729 [1983]), as well as within the coding sequence itself (Osborne et al., Mol. Cell Bio. 4:1293 [1984]). Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin,  $\alpha$ -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, Nature 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the Htk ligand-encoding sequence, but is preferably located at a site 5' from the promoter.

(vi) Transcription Termination Component

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding Htk ligand.

(vii) Construction and Analysis of Vectors

Construction of suitable vectors containing one or more of the above listed components employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required.

- 5 For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to transform *E. coli* K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequenced  
10 by the method of Messing et al., Nucleic Acids Res. 9:309 (1981) or by the method of Maxam et al., Methods in Enzymology 65:499 (1980).

(viii) Transient Expression Vectors

- Particularly useful in the practice of this invention are expression vectors that provide for the transient expression in mammalian cells of DNA  
15 encoding Htk ligand. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector. Sambrook et al., *supra*, pp. 16.17 - 16.22.  
20 Transient expression systems, comprising a suitable expression vector and a host cell, allow for the convenient positive identification of polypeptides encoded by cloned DNAs, as well as for the rapid screening of such polypeptides for desired biological or physiological properties. Thus, transient expression systems are particularly useful in the invention  
25 for purposes of identifying analogs and variants of Htk ligand that are biologically active Htk ligand.

(ix) Suitable Exemplary Vertebrate Cell Vectors

- Other methods, vectors, and host cells suitable for adaptation to the synthesis of Htk ligand in recombinant vertebrate cell culture are  
30 described in Gething et al., Nature 293:620-625 (1981); Mantei et al., Nature 281:40-46 (1979); Levinson et al.; EP 117,060; and EP 117,058. A particularly useful plasmid for mammalian cell culture expression of Htk ligand is pRK5 (EP 307,247) or pSVI6B (PCT pub. no. WO 91/08291 published 13 June 1991).

35 D. Selection and Transformation of Host Cells

- Suitable host cells for cloning or expressing the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as  
40 *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. One  
45 preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110

(ATCC 27,325) are suitable. These examples are illustrative rather than limiting. Strain W3110 is a particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell should secrete minimal amounts  
 5 of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins, with examples of such hosts including *E. coli* W3110 strain 27C7. The complete genotype of 27C7 is *tonAΔ ptr3 phoAΔE15 Δ(argF-lac)169 ompTΔ degP41karf*. Strain 27C7 was deposited on 30 October 1991 in the American Type Culture Collection  
 10 as ATCC No. 55,244. Alternatively, the strain of *E. coli* having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990 may be employed. Alternatively, methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous  
 15 fungi or yeast are suitable cloning or expression hosts for Htk ligand-encoding vectors. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe* (Beach and  
 20 Nurse, Nature 290:140 [1981]; EP 139,383 published May 2, 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., *supra*) such as, e.g., *K. lactis* [MW98-8C, CBS683, CBS4574; Louvencourt et al., J. Bacteriol., 737 (1983)], *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K.*  
 25 *drosophilorum* (ATCC 36,906; Van den Berg et al., *supra*), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* [EP 402,226]; *Pichia pastoris* (EP 183,070; Sreekrishna et al., J. Basic Microbiol. 28:265-278 [1988]); *Candida*; *Trichoderma reesia* [EP 244,234]; *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA 76:5259-5263 [1979]); *Schwanniomyces* such as  
 30 *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun. 112:284-289 [1983]; Tilburn et al., Gene 26:205-221 [1983]; Yelton et al., Proc. Natl.  
 35 Acad. Sci. USA 81:1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J. 4:475-479 [1985]).

Suitable host cells for the expression of glycosylated Htk ligand are derived from multicellular organisms. Such host cells are capable of complex processing and glycosylation activities. In principle, any higher  
 40 eukaryotic cell culture is workable, whether from vertebrate or invertebrate culture. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito),  
 45 *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. See, e.g., Luckow et al., Bio/Technology 6:47-55 (1988); Miller et al., in

Genetic Engineering, Setlow et al., eds., Vol. 8 (Plenum Publishing, 1986), pp. 277-279; and Maeda et al., Nature 315:592-594 (1985). A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can be utilized as hosts. Typically, plant cells are transfected by incubation with certain strains of the bacterium *Agrobacterium tumefaciens*, which has been previously manipulated to contain the Htk ligand DNA. During incubation of the plant cell culture with *A. tumefaciens*, the DNA encoding the Htk ligand is transferred to the plant cell host such that it is transfected, and will, under appropriate conditions, express the Htk ligand DNA. In addition, regulatory and signal sequences compatible with plant cells are available, such as the nopaline synthase promoter and polyadenylation signal sequences. Depicker et al., J. Mol. Appl. Gen. 1:561 (1982). In addition, DNA segments isolated from the upstream region of the T-DNA 780 gene are capable of activating or increasing transcription levels of plant-expressible genes in recombinant DNA-containing plant tissue. EP 321,196 published 21 June 1989.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure in recent years (Tissue Culture, Academic Press, Kruse and Patterson, editors [1973]). Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol. 36:59 [1977]); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA 77:4216 [1980]); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 [1980]); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad. Sci. 383:44-68 [1982]); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

Host cells are transfected and preferably transformed with the above-described expression or cloning vectors of this invention and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

Transfection refers to the taking up of an expression vector by a host cell whether or not any coding sequences are in fact expressed. Numerous methods of transfection are known to the ordinarily skilled

artisan, for example,  $\text{CaPO}_4$  and electroporation. Successful transfection is generally recognized when any indication of the operation of this vector occurs within the host cell.

Transformation means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook et al., *supra*, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene 23:315 (1983) and WO 89/05859 published 29 June 1989. In addition, plants may be transfected using ultrasound treatment as described in WO 91/00358 published 10 January 1991.

For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology 52:456-457 (1978) is preferred. General aspects of mammalian cell host system transformations have been described by Axel in U.S. Patent No. 4,399,216 issued 16 August 1983. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact. 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA) 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, etc., may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology (1989), Keown et al., Methods in Enzymology 185:527-537 (1990), and Mansour et al., Nature 336:348-352 (1988).

#### E. Culturing the Host Cells

Prokaryotic cells used to produce the Htk ligand polypeptide of this invention are cultured in suitable media as described generally in Sambrook et al., *supra*.

The mammalian host cells used to produce the Htk ligand of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ([MEM], Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ([DMEM], Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham and Wallace, Meth. Enz. 58:44 (1979), Barnes and Sato, Anal. Biochem. 102:255 (1980), U.S. Patent Nos. 4,767,704; 4,657,866; 4,927,762; or 4,560,655; WO 90/03430; WO 87/00195; U.S. Patent Re. 30,985; or U.S. Patent No. 5,122,469, the disclosures of all of which are incorporated herein by reference, may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and



thymidine), antibiotics (such as Gentamycin<sup>TM</sup> drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

In general, principles, protocols, and practical techniques for maximizing the productivity of mammalian cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed., IRL Press, 1991.

The host cells referred to in this disclosure encompass cells in culture as well as cells that are within a host animal.

15 F. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA 77:5201-5205 [1980]), dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Various labels may be employed, most commonly radioisotopes, particularly <sup>32</sup>P. However, other techniques may also be employed, such as using biotin-modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. With immunohistochemical staining techniques, a cell sample is prepared, typically by dehydration and fixation, followed by reaction with labeled antibodies specific for the gene product coupled, where the labels are usually visually detectable, such as enzymatic labels, fluorescent labels, luminescent labels, and the like. A particularly sensitive staining technique suitable for use in the present invention is described by Hsu et al., Am. J. Clin. Path. 75:734-738 (1980).

Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a

native Htk ligand polypeptide or against a synthetic peptide based on the DNA sequences provided herein as described further in Section 4 below.

G. Purification of Htk Ligand Polypeptide

Htk ligand preferably is recovered from the culture medium as a secreted polypeptide, although it also may be recovered from host cell lysates when directly produced without a secretory signal. If the Htk ligand is membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100)

When Htk ligand is produced in a recombinant cell other than one of human origin, the Htk ligand is completely free of proteins or polypeptides of human origin. However, it is necessary to purify Htk ligand from recombinant cell proteins or polypeptides to obtain preparations that are substantially homogeneous as to Htk ligand. As a first step, the culture medium or lysate is centrifuged to remove particulate cell debris. Htk ligand thereafter is purified from contaminant soluble proteins and polypeptides, with the following procedures being exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; and protein A Sepharose columns to remove contaminants such as IgG.

In the preferred embodiment, the Htk receptor-Fc fusion disclosed in Bennett et al., *supra*, is immobilized on a protein A Sepharose column and the Htk ligand can be isolated by affinity purification using this column.

Htk ligand variants in which residues have been deleted, inserted, or substituted are recovered in the same fashion as native Htk ligand, taking account of any substantial changes in properties occasioned by the variation. For example, preparation of a Htk ligand fusion with another protein or polypeptide, e.g., a bacterial or viral antigen, facilitates purification; an immunoaffinity column containing antibody to the antigen can be used to adsorb the fusion polypeptide. Immunoaffinity columns such as a rabbit polyclonal anti-Htk ligand column can be employed to absorb the Htk ligand variant by binding it to at least one remaining immune epitope. A protease inhibitor such as phenyl methyl sulfonyl fluoride (PMSF) also may be useful to inhibit proteolytic degradation during purification, and antibiotics may be included to prevent the growth of adventitious contaminants. One skilled in the art will appreciate that purification methods suitable for native Htk ligand may require modification to account for changes in the character of Htk ligand or its variants upon expression in recombinant cell culture.

H. Covalent Modifications of Htk Ligand Polypeptides

Covalent modifications of Htk ligand polypeptides are included within the scope of this invention. Both native Htk ligand and amino acid sequence variants of the Htk ligand may be covalently modified. One type of covalent modification included within the scope of this invention is a Htk ligand fragment (e.g. soluble Htk ligand). Variant Htk ligand

fragments having up to about 40 amino acid residues may be conveniently prepared by chemical synthesis or by enzymatic or chemical cleavage of the full-length or variant Htk ligand polypeptide. Other types of covalent modifications of the Htk ligand or fragments thereof are introduced into the molecule by reacting targeted amino acid residues of the Htk ligand or fragments thereof with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

Cysteiny l residues most commonly are reacted with  $\alpha$ -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteiny l residues also are derivatized by reaction with bromotrifluoroacetone,  $\alpha$ -bromo- $\beta$ -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

Lysiny l and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysiny l residues. Other suitable reagents for derivatizing  $\alpha$ -amino-containing residues include imidoesters such as methyl picolinimate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginy l residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high  $pK_a$  of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using  $^{125}I$  or  $^{131}I$  to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ( $R-N=C=N-R'$ ), where R and R' are different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl)carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy l and glutaminy l residues by reaction with ammonium ions.

Derivatization with bifunctional agents is useful for crosslinking Htk ligand to a water-insoluble support matrix or surface for use in the method for purifying anti-Htk ligand antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates described in U.S. Patent Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

Glutaminy and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. These residues are deamidated under neutral or basic conditions. The deamidated form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 [1983]), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the Htk ligand polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. By altering is meant deleting one or more carbohydrate moieties found in native Htk ligand, and/or adding one or more glycosylation sites that are not present in the native Htk ligand.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-aceylgalactosamine, galactose, or xylose to a hydroxylamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the Htk ligand polypeptide is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the

addition of, or substitution by, one or more serine or threonine residues to the native Htk ligand sequence (for O-linked glycosylation sites). For ease, the Htk ligand amino acid sequence is preferably altered through changes at the DNA level, particularly by mutating the DNA encoding the Htk ligand polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids. The DNA mutation(s) may be made using methods described above under the heading of "Amino Acid Sequence Variants of Native Htk Ligand."

Another means of increasing the number of carbohydrate moieties on the Htk ligand polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. These procedures are advantageous in that they do not require production of the polypeptide in a host cell that has glycosylation capabilities for N- or O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the Htk ligand polypeptide may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the polypeptide to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin, et al., Arch. Biochem. Biophys. 259:52 (1987) and by Edge et al., Anal. Biochem. 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., Meth. Enzymol. 138:350 (1987).

Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., J. Biol. Chem. 257:3105 (1982). Tunicamycin blocks the formation of protein-N-glycoside linkages.

Another type of covalent modification of Htk ligand comprises linking the Htk ligand polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

Since it is often difficult to predict in advance the characteristics of a variant Htk ligand, it will be appreciated that some screening of the recovered variant will be needed to select the optimal variant. A change in the immunological character of the Htk ligand molecule, such as affinity for a given antibody, is also able to be measured by a competitive-type

immunoassay. The variant is assayed for changes in the suppression or enhancement of its enzymatic activity by comparison to the activity observed for native Htk ligand in the same assay. For example, one can screen for the ability of the variant Htk ligand to stimulate protein kinase activity of the Htk receptor using the techniques set forth in Lokker et al., EMBO 11:2503-2510 (1992). See also Example 4 herein. Other potential modifications of protein or polypeptide properties such as redox or thermal stability, hydrophobicity, susceptibility to proteolytic degradation, or the tendency to aggregate with carriers or into multimers are assayed by methods well known in the art.

#### I. Htk Ligand-ImmunoglobulinChimeras (Immunoadhesins)

Immunoglobulins (Ig) and certain variants thereof are known and many have been prepared in recombinant cell culture. For example, see U.S. Patent No. 4,745,055; EP 256,654; Faulkner et al., Nature 298:286 (1982); EP 120,694; EP 125,023; Morrison, J. Immun. 123:793 (1979); Köhler et al., Proc. Natl. Acad. Sci. USA 77:2197 (1980); Raso et al., Cancer Res. 41:2073 (1981); Morrison et al., Ann. Rev. Immunol. 2:239 (1984); Morrison, Science 229:1202 (1985); Morrison et al., Proc. Natl. Acad. Sci. USA 81:6851 (1984); EP 255,694; EP 266,663; and WO 88/03559. Reassorted immunoglobulin chains also are known. See, for example, U.S. Patent No. 4,444,878; WO 88/03565; and EP 68,763 and references cited therein. Chimeras constructed from a receptor sequence linked to an appropriate immunoglobulin constant domain sequence (immunoadhesins) are known in the art. Immunoadhesins reported in the literature include fusions of the T cell receptor (Gascoigne et al., Proc. Natl. Acad. Sci. USA 84:2936-2940 [1987]); CD4 (Capon et al., Nature 337:525-531 [1989]; Traunecker et al., Nature 339:68-70 [1989]; Zettmeissl et al., DNA Cell Biol. USA 9:347-353 [1990]; and Byrn et al., Nature 344:667-670 [1990]); L-selectin (homing receptor) (Watson et al., J. Cell. Biol. 110:2221-2229 [1990]; and Watson et al., Nature 349:164-167 [1991]); CD44 (Aruffo et al., Cell 61:1303-1313 [1990]); CD28 and B7 (Linsley et al., J. Exp. Med. 173:721-730 [1991]); CTLA-4 (Lisley et al., J. Exp. Med. 174:561-569 [1991]); CD22 (Stamenkovic et al., Cell 66:1133-1144 [1991]); TNF receptor (Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 [1991]; Lesslauer et al., Eur. J. Immunol. 27:2883-2886 [1991]; and Peppel et al., J. Exp. Med. 174:1483-1489 [1991]); and IgE receptor  $\alpha$  (Ridgway and Gorman, J. Cell. Biol. Vol. 115, abstr. 1448 [1991]).

The simplest and most straightforward immunoadhesin design combined the binding region(s) of the "adhesin" protein with the hinge and Fc regions of an immunoglobulin heavy chain. Ordinarily, when preparing the Htk ligand-immunoglobulin chimeras of the present invention, nucleic acid encoding the extracellular domain of the Htk ligand, or a fragment thereof, will be fused C-terminally to nucleic acid encoding the N-terminus of an immunoglobulin constant domain sequence, however N-terminal fusions are also possible.

Typically, in such fusions the encoded chimeric polypeptide will retain at least functionally active hinge, CH2 and CH3 domains of the constant region of an immunoglobulin heavy chain. Fusions are also made to the C-terminus of the Fc portion of a constant domain, or immediately  
 5 N-terminal to the CH1 of the heavy chain or the corresponding region of the light chain.

The precise site at which the fusion is made is not critical; particular sites are well known and may be selected in order to optimize the biological activity, secretion, or binding characteristics of the Htk  
 10 ligand-immunoglobulin chimeras.

In some embodiments, the Htk ligand-immunoglobulin chimeras are assembled as monomers, or hetero- or homo-multimers, and particularly as dimers or tetramers, essentially as illustrated in WO 91/08298.

In a preferred embodiment, the Htk ligand extracellular domain  
 15 sequence is fused to the N-terminus of the Fc domain of immunoglobulin G<sub>1</sub> (IgG-1). It is possible to fuse the entire heavy chain constant region to the Htk ligand extracellular domain sequence. However, more preferably, a sequence beginning in the hinge region just upstream of the papain cleavage site which defines IgG Fc chemically (i.e. residue 216, taking the  
 20 first residue of heavy chain constant region to be 114), or analogous sites of other immunoglobulins is used in the fusion. In a particularly preferred embodiment, the Htk ligand amino acid sequence is fused to (a) the hinge region and CH2 and CH3 or (b) the CH1, hinge, CH2 and CH3 domains, of an IgG-1, IgG-2, or IgG-3 heavy chain. The precise site at  
 25 which the fusion is made is not critical, and the optimal site can be determined by routine experimentation.

In some embodiments, the Htk ligand-immunoglobulin chimeras are assembled as multimers, and particularly as homo-dimers or -tetramers. Generally, these assembled immunoglobulins will have known unit structures.  
 30 A basic four chain structural unit is the form in which IgG, IgD, and IgE exist. A four chain unit is repeated in the higher molecular weight immunoglobulins; IgM generally exists as a pentamer of four basic units held together by disulfide bonds. IgA globulin, and occasionally IgG globulin, may also exist in multimeric form in serum. In the case of  
 35 multimer, each of the four units may be the same or different.

Various exemplary assembled Htk ligand-immunoglobulin chimeras within the scope herein are schematically diagrammed below:

- (a)  $AC_L-AC_H$ ;
- (b)  $AC_H-[AC_H, AC_L-AC_H, AC_L-V_HC_H, \text{ or } V_LC_L-AC_H]$ ;
- 40 (c)  $AC_L-AC_H-[AC_L-AC_H, AC_L-V_HC_H, V_LC_L-AC_H, \text{ or } V_LC_L-V_HC_H]$ ;
- (d)  $AC_L-V_HC_H-[AC_H, \text{ or } AC_L-V_HC_H, \text{ or } V_LC_L-AC_H]$ ;
- (e)  $V_LC_L-AC_H-[AC_L-V_HC_H, \text{ or } V_LC_L-AC_H]$ ; and
- (f)  $[A-Y]_n-[V_LC_L-V_HC_H]_n$ ,

wherein

45 each A represents identical or different Htk ligand amino acid sequences;

V<sub>L</sub> is an immunoglobulin light chain variable domain;  
V<sub>H</sub> is an immunoglobulin heavy chain variable domain;  
C<sub>L</sub> is an immunoglobulin light chain constant domain;  
C<sub>H</sub> is an immunoglobulin heavy chain constant domain;

5 n is an integer greater than 1;

Y designates the residue of a covalent cross-linking agent.

In the interests of brevity, the foregoing structures only show key features; they do not indicate joining (J) or other domains of the immunoglobulins, nor are disulfide bonds shown. However, where such domains are required for binding activity, they shall be constructed to be present in the ordinary locations which they occupy in the immunoglobulin molecules.

Alternatively, the Htk ligand extracellular domain sequences can be inserted between immunoglobulin heavy chain and light chain sequences, such that an immunoglobulin comprising a chimeric heavy chain is obtained. In this embodiment, the Htk ligand sequences are fused to the 3' end of an immunoglobulin heavy chain in each arm of an immunoglobulin, either between the hinge and the CH2 domain, or between the CH2 and CH3 domains. Similar constructs have been reported by Hoogenboom, et al., Mol. Immunol. 28:1027-1037 (1991).

Although the presence of an immunoglobulin light chain is not required in the immunoadhesins of the present invention, an immunoglobulin light chain might be present either covalently associated to an Htk ligand-immunoglobulin heavy chain fusion polypeptide, or directly fused to the Htk ligand extracellular domain. In the former case, DNA encoding an immunoglobulin light chain is typically coexpressed with the DNA encoding the Htk ligand-immunoglobulin heavy chain fusion protein. Upon secretion, the hybrid heavy chain and the light chain will be covalently associated to provide an immunoglobulin-like structure comprising two disulfide-linked immunoglobulin heavy chain-light chain pairs. Methods suitable for the preparation of such structures are, for example, disclosed in U.S. Patent No. 4,816,567, issued 28 March 1989.

In a preferred embodiment, the immunoglobulin sequences used in the construction of the immunoadhesins of the present invention are from an IgG immunoglobulin heavy chain constant domain. For human immunoadhesins, the use of human IgG1 and IgG3 immunoglobulin sequences is preferred. A major advantage of using IgG1 is that IgG1 immunoadhesins can be purified efficiently on immobilized protein A. In contrast, purification of IgG3 requires protein G, a significantly less versatile medium. However, other structural and functional properties of immunoglobulins should be considered when choosing the Ig fusion partner for a particular immunoadhesin construction. For example, the IgG3 hinge is longer and more flexible, so it can accommodate larger "adhesin" domains that may not fold or function properly when fused to IgG1. Another consideration may be valency; IgG immunoadhesins are bivalent homodimers, whereas Ig subtypes like IgA and IgM may give rise to dimeric or pentameric structures,



respectively, of the basic Ig homodimer unit. For Htk ligand-Ig immunoadhesins designed for in vivo application, the pharmacokinetic properties and the effector functions specified by the Fc region are important as well. Although IgG1, IgG2 and IgG4 all have in vivo half-lives of 21 days, their relative potencies at activating the complement system are different. IgG4 does not activate complement, and IgG2 is significantly weaker at complement activation than IgG1. Moreover, unlike IgG1, IgG2 does not bind to Fc receptors on mononuclear cells or neutrophils. While IgG3 is optimal for complement activation, its in vivo half-life is approximately one third of the other IgG isotypes. Another important consideration for immunoadhesins designed to be used as human therapeutics is the number of allotypic variants of the particular isotype. In general, IgG isotypes with fewer serologically-defined allotypes are preferred. For example, IgG1 has only four serologically-defined allotypic sites, two of which (G1m and 2) are located in the Fc region; and one of these sites, G1m1, is non-immunogenic. In contrast, there are 12 serologically-defined allotypes in IgG3, all of which are in the Fc region; only three of these sites (G3m5, 11 and 21) have one allotype which is nonimmunogenic. Thus, the potential immunogenicity of a  $\gamma$ 3 immunoadhesin is greater than that of a  $\gamma$ 1 immunoadhesin.

In designing the Htk ligand-Ig immunoadhesins of the present invention, domains that are not required for rPTK binding and/or biological activity of the Htk ligand may be deleted. With respect to the parental immunoglobulin, a useful joining point is just upstream of the cysteines of the hinge that form the disulfide bonds between the two heavy chains. In a frequently used design, the codon for the C-terminal residue of the "adhesin" (Htk ligand) part of the molecule is placed directly upstream of the codons for the sequence DKHTCTCPPCP (SEQ ID NO: 7) of the IgG1 hinge region.

The general methods suitable for the construction and expression of immunoadhesins are the same as those disclosed hereinabove with regard to (native or variant) Htk ligand. Htk ligand-Ig immunoadhesins are most conveniently constructed by fusing the cDNA sequence encoding the Htk ligand portion in-frame to an Ig cDNA sequence. However, fusion to genomic Ig fragments can also be used (see, e.g. Gascoigne et al., *supra*; Aruffo et al., *Cell* 61:1303-1313 [1990]; and Stamenkovic et al., *Cell* 66:1133-1144 [1991]). The latter type of fusion requires the presence of Ig regulatory sequences for expression. cDNAs encoding IgG heavy-chain constant regions can be isolated based on published sequences from cDNA libraries derived from spleen or peripheral blood lymphocytes, by hybridization or by polymerase chain reaction (PCR) techniques. The cDNAs encoding the "adhesin" and the Ig parts of the immunoadhesin are inserted in tandem into a plasmid vector that directs efficient expression in the chosen host cells. For expression in mammalian cells, pRK5-based vectors (Schall et al., *Cell* 61:361-370 [1990]) and CDM8-based vectors (Seed, *Nature* 329:840 [1989]) are useful. The exact junction can be created by removing the

extra sequences between the designed junction codons using oligonucleotide-directed deletional mutagenesis (Zoller and Smith, Nucleic Acids Res. 10:6487 [1982]; and Capon et al., Nature 337:525-531 [1989]). Synthetic oligonucleotides can be used, in which each half is complementary to the  
5 sequence on either side of the desired junction; ideally, these are 36- to 48-mers. Alternatively, PCR techniques can be used to join the two parts of the molecule in-frame with an appropriate vector.

The choice of host cell line for the expression of Htk ligand-Ig immunoadhesins depends mainly on the expression vector. Another  
10 consideration is the amount of protein that is required. Milligram quantities often can be produced by transient transfections. For example, the adenovirus EIA-transformed 293 human embryonic kidney cell line can be transfected transiently with pRK5-based vectors by a modification of the calcium phosphate method to allow efficient immunoadhesin expression.  
15 CDM8-based vectors can be used to transfect COS cells by the DEAE-dextran method (Aruffo et al., Cell 61:1303-1313 [1990]; and Zettmeissl et al., DNA Cell Biol. (US) 9:347-353 [1990]). If larger amounts of protein are desired, the immunoadhesin can be expressed after stable transfection of a host cell line. For example, a pRK5-based vector can be introduced into  
20 Chinese hamster ovary (CHO) cells in the presence of an additional plasmid encoding dihydrofolate reductase (DHFR) and conferring resistance to G418. Clones resistant to G418 can be selected in culture. These clones are grown in the presence of increasing levels of DHFR inhibitor methotrexate and clones are selected in which the number of gene copies encoding the DHFR  
25 and immunoadhesin sequences is co-amplified. If the immunoadhesin contains a hydrophobic leader sequence at its N-terminus, it is likely to be processed and secreted by the transfected cells. The expression of immunoadhesins with more complex structures may require uniquely suited host cells. For example, components such as light chain or J chain may be  
30 provided by certain myeloma or hybridoma host cells (Gascoigne et al., *supra*; and Martin et al., J. Virol. 67:3561-3568 [1993]).

Immunoadhesins can be conveniently purified by affinity chromatography. The suitability of protein A as an affinity ligand depends on the species and isotype of the immunoglobulin Fc domain that is used in  
35 the chimera. Protein A can be used to purify immunoadhesins that are based on human  $\gamma 1$ ,  $\gamma 2$ , or  $\gamma 4$  heavy chains (Lindmark et al., J. Immunol. Meth. 62:1-13 [1983]). Protein G is recommended for all mouse isotypes and for human  $\gamma 3$  (Guss et al., EMBO J. 5:1567-1575 [1986]). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are  
40 available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. The conditions for binding an immunoadhesin to the protein A or G affinity column are dictated entirely by the characteristics of the Fc domain; that is, its species and  
45 isotype. Generally, when the proper ligand is chosen, efficient binding occurs directly from unconditioned culture fluid. One distinguishing

feature of immunoadhesins is that, for human  $\gamma 1$  molecules, the binding capacity for protein A is somewhat diminished relative to an antibody of the same Fc type. Bound immunoadhesin can be efficiently eluted either at acidic pH (at or above 3.0), or in a neutral pH buffer containing a mildly chaotropic salt. This affinity chromatography step can result in an immunoadhesin preparation that is >95% pure.

Other methods known in the art can be used in place of, or in addition to, affinity chromatography on protein A or G to purify immunoadhesins. Immunoadhesins behave similarly to antibodies in thiophilic gel chromatography (Hutchens and Porath, Anal. Biochem. 159:217-226 [1986]) and immobilized metal chelate chromatography (Al-Mashikhi and Makai, J. Dairy Sci. 71:1756-1763 [1988]). In contrast to antibodies, however, their behavior on ion exchange columns is dictated not only by their isoelectric points, but also by a charge dipole that may exist in the molecules due to their chimeric nature.

#### J. Epitope tagged Htk Ligand

This application encompasses chimeric polypeptides comprising Htk ligand fused to another polypeptide (such as the immunoadhesins mentioned above). In one preferred embodiment, the chimeric polypeptide comprises a fusion of the Htk ligand (or a fragment thereof, e.g., the ECD of the Htk ligand) with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally provided at the amino- or carboxyl- terminus of the Htk ligand. Such epitope tagged forms of the Htk ligand are desirable, as the presence thereof can be detected using a labelled antibody against the tag polypeptide. Also, provision of the epitope tag enables the Htk ligand to be readily purified by affinity purification using the anti-tag antibody. Affinity purification techniques and diagnostic assays involving antibodies are described later herein.

Tag polypeptides and their respective antibodies are well known in the art. Examples include the flu HA tag polypeptide and its antibody 12CA5, (Field et al., Mol. Cell. Biol. 8:2159-2165 [1988]); the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto (Evan et al., Molecular and Cellular Biology 5(12):3610-3616 [1985]); and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody (Paborsky et al., Protein Engineering 3(6):547-553 [1990]). Other tag polypeptides have been disclosed. Examples include the Flag-peptide (Hopp et al., BioTechnology 6:1204-1210 [1988]); the KT3 epitope peptide (Martin et al., Science 255:192-194 [1992]); an  $\alpha$ -tubulin epitope peptide (Skinner et al., J. Biol. Chem 266:15163-15166 [1991]); and the T7 gene 10 protein peptide tag (Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA 87:6393-6397 [1990]). Once the tag polypeptide has been selected, an antibody thereto can be generated using the techniques disclosed herein.

The general methods suitable for the construction and production of epitope tagged Htk ligand are the same as those disclosed hereinabove with regard to (native or variant) Htk ligand. Htk ligand-tag polypeptide

fusions are most conveniently constructed by fusing the cDNA sequence encoding the Htk ligand portion in-frame to the tag polypeptide DNA sequence and expressing the resultant DNA fusion construct in appropriate host cells. Ordinarily, when preparing the Htk ligand-tag polypeptide  
5 chimeras of the present invention, nucleic acid encoding the Htk ligand (or a fragment thereof) will be fused at its 3' end to nucleic acid encoding the N-terminus of the tag polypeptide, however 5' fusions are also possible.

Epitope tagged Htk ligand can be conveniently purified by affinity  
10 chromatography using the anti-tag antibody. The matrix to which the affinity antibody is attached is most often agarose, but other matrices are available (e.g. controlled pore glass or poly(styrenedivinyl)benzene). The epitope tagged Htk ligand can be eluted from the affinity column by varying the buffer pH or ionic strength or adding chaotropic agents, for example.

15 2. Therapeutic Uses, Compositions and Administration of Htk Ligand

Htk ligand is believed to find therapeutic use for treating mammals via stimulation or inhibition of growth and/or differentiation and/or activation of cells having a receptor for the Htk ligand, such as the Htk receptor. The prominent regional expression of Htk ligand DNA in the  
20 cerebral cortex, hippocampus, striatum, and cerebellum (see Example 3) suggests the possibility that Htk ligand polypeptide might be useful to treat neurodegenerative diseases in which these structures, or neurons projecting to these structures, are affected. Such diseases include, but are not limited to, Alzheimer's disease, Parkinson's disease, Huntington's  
25 chorea, and disorders of the cerebellum (Hefti, J. Neurobiol. in press [1994]; Marsden, Lancet 335:948-952 [1990]; Agid, Lancet 337:1321-1327 [1991]; Wexler et al., Ann. Rev. Neurosci. 14:503-529 [1991]).

Mature exogenous Htk ligand or a soluble form thereof (e.g., a soluble immunoadhesin) may be administered to a patient in these  
30 circumstances. The human Htk ligand is clearly useful insofar as it can be administered to a human having depressed levels of endogenous Htk ligand, preferably in the situation where such depressed levels lead to a pathological disorder.

Therapeutic formulations of Htk ligand are prepared for storage by  
35 mixing Htk ligand having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (Remington's Pharmaceutical Sciences, 16th edition, Osol, A., Ed., (1980)), in the form of lyophilized cake or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and  
40 concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine,  
45 asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents

such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, Pluronic or polyethylene glycol (PEG).

Htk ligand also may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-[methacrylate] microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, supra.

Htk ligand to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. Htk ligand ordinarily will be stored in lyophilized form or in solution.

Therapeutic Htk ligand compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The route of Htk ligand administration is in accord with known methods, e.g., injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial, or intralesional routes, or by sustained release systems as noted below. Htk ligand is administered continuously by infusion or by bolus injection. Htk ligand antibody is administered in the same fashion, or by administration into the blood stream or lymph.

Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the protein, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels [e.g., poly(2-hydroxyethyl-methacrylate) as described by Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981) and Langer, Chem. Tech. 12:98-105 (1982) or poly(vinylalcohol)], polylactides (U.S. Patent No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al., Biopolymers 22:547-556 [1983]), non-degradable ethylene-vinyl acetate (Langer et al., supra), degradable lactic acid-glycolic acid copolymers such as the Lupron Depot™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated proteins remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for protein stabilization depending on the

mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Sustained-release Htk ligand compositions also include liposomally entrapped Htk ligand. Liposomes containing Htk ligand are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. USA 82:3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. Patent Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the optimal Htk ligand therapy.

An effective amount of Htk ligand to be employed therapeutically will depend, for example, upon the therapeutic objectives, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage might range from about 1 µg/kg to up to 10 mg/kg or more, depending on the factors mentioned above. Typically, the clinician will administer Htk ligand until a dosage is reached that achieves the desired effect. The progress of this therapy is easily monitored by conventional assays.

### 3. Non-Therapeutic, Diagnostic Uses for Htk Ligand

The nucleic acid encoding the Htk ligand may be used as a diagnostic for tissue-specific typing (e.g. mammary gland epithelia). For example, such procedures as *in situ* hybridization, Northern and Southern blotting, and PCR analysis may be used to determine whether DNA and/or RNA encoding Htk ligand is present in the cell type(s) being evaluated. Htk ligand nucleic acid or polypeptide may also be used as diagnostic markers for mammary gland carcinomas. For example, the Htk ligand may be labeled, using the techniques described herein, and expression of Htk receptor nucleic acid can be quantified, using the labelled Htk ligand.

Human Htk ligand nucleic acid has been localized to chromosome 13q33. Thus, the nucleic acid for human Htk ligand can be used as a marker for this human chromosome.

Htk ligand nucleic acid is also useful for the preparation of Htk ligand polypeptide by recombinant techniques exemplified herein.

Isolated Htk ligand polypeptide may be used in quantitative diagnostic assays as a standard or control against which samples containing unknown quantities of Htk ligand may be prepared.

Htk ligand preparations are also useful in generating antibodies, as standards in assays for Htk ligand (e.g., by labeling Htk ligand for use as a standard in a radioimmunoassay, or enzyme-linked immunoassay), for detecting the presence of the Htk receptor in a biological sample (e.g.,  
5 using a labelled Htk ligand) in affinity purification techniques, and in competitive-type receptor binding assays when labeled with radioiodine, enzymes, fluorophores, spin labels, and the like.

The Htk ligand is also useful as a diagnostic tool. For example, the Htk ligand can be produced in prokaryotic cells using the techniques  
10 elaborated herein and the unglycosylated protein so produced can be used as a molecular weight marker, for example. The deduced molecular weight (mw) of the unglycosylated Htk ligand under reducing conditions is about 34 kD. Soluble Htk ligand has a deduced mw of 22 kD under reducing conditions. In order to use Htk ligand as a molecular weight marker, gel  
15 filtration chromatography or SDS-PAGE, for example, will be used to separate protein(s) for which it is desired to determine their molecular weight(s) in substantially the normal way. The Htk ligand and other molecular weight markers will be used as standards to provide a range of molecular weights. For example, phosphorylase b (mw = 97,400), bovine serum  
20 albumin (mw = 68,000), ovalbumin (mw = 46,000), Htk ligand (mw = 34,000), trypsin inhibitor (mw = 20,100), and lysozyme (mw = 14,400) can be used as mw markers. The other molecular weight markers mentioned here can be purchased commercially from Amersham Corporation, Arlington Heights, IL, for example. Often, the molecular weight markers will be labelled to enable  
25 easy detection following separation. Techniques for labelling antibodies and proteins are discussed herein and are well known in the art. For example, the molecular weight markers may be biotinylated and, following separation on SDS-PAGE, for example, the blot can be incubated with streptavidin-horseradish peroxidase. The bands can then be detected by  
30 light detection.

It may also be useful to grow certain cells having the Htk receptor ex vivo using the Htk ligand as a growth factor. These cells which are to be grown ex vivo may simultaneously be exposed to other known growth factors or cytokines. Exemplary cytokines include the interleukins (e.g.,  
35 IL-3), granulocyte-macrophage colony-stimulating factor (GM-CSF), macrophage colony-stimulating factor (M-CSF), granulocyte colony-stimulating factor (GM-CSF), erythropoietin (Epo), lymphotoxin, steel factor (SLF), tumor necrosis factor (TNF) and gamma-interferon. This results in proliferation and/or differentiation of the cells having the Htk  
40 receptor. For example, human tumor cell lines for which it is desired to isolate certain tumor associated factors (usually proteins) therefrom may be grown ex vivo using the Htk ligand. Also, antibodies against the tumor associated factors can be generated which may be useful for diagnostic purposes. Examples of such tumor cell lines which can be treated with the  
45 Htk ligand include mammary cancer cells (e.g. MCF-7), liver cell lines, Colo 205, NCI 69, HM-1 and HeLa, for example.

In yet another aspect of the invention, the ligand may be used for affinity purification of the Htk receptor. Briefly, this technique involves covalently attaching the Htk ligand to an inert and porous matrix (e.g., agarose reacted with cyanogen bromide). A solution containing the  
5 Htk receptor can then be passed through the chromatographic material and can be subsequently released by changing the elution conditions (e.g. by changing pH or ionic strength).

The purified Htk ligand, and the nucleic acid encoding it, may also be sold as reagents for mechanism studies of the ligand and its cognate  
10 receptor, to study the role of the Htk ligand and receptor in normal growth and development, as well as abnormal growth and development, e.g. in malignancies.

Htk ligand may be used for competitive screening of potential agonists or antagonists for binding to the Htk receptor. Htk ligand  
15 variants are useful as standards or controls in assays for Htk ligand, provided that they are recognized by the analytical system employed, e.g. an anti-Htk ligand antibody.

#### 4. Htk ligand Antibody Preparation

A description follows as to the production of exemplary antibodies  
20 as defined herein. These exemplary antibodies include polyclonal, monoclonal, humanized, bispecific or heteroconjugate antibodies.

##### A. Polyclonal Antibodies

Polyclonal antibodies to the Htk ligand generally are raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections  
25 of the Htk ligand and an adjuvant. It may be useful to conjugate the Htk ligand or a fragment containing the target amino acid sequence to a protein that is immunogenic in the species to be immunized, e.g., keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example  
30 maleimidobenzoyl sulfo succinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glytaraldehyde, succinic anhydride,  $\text{SOCl}_2$ , or  $\text{R}^1\text{N}=\text{C}=\text{NR}$ , where R and  $\text{R}^1$  are different alkyl groups.

Animals are immunized against the immunogenic conjugates or  
35 derivatives by combining 1 mg of 1  $\mu\text{g}$  of conjugate (for rabbits or mice, respectively) with 3 volumes of Freud's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of conjugate in Freud's complete adjuvant by subcutaneous injection at multiple sites. 7 to 14  
40 days later the animals are bled and the serum is assayed for anti-Htk ligand antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same Htk ligand, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell



culture as protein fusions. Also, aggregating agents such as alum are used to enhance the immune response.

#### B. Monoclonal Antibodies

Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

For example, the anti-Htk ligand monoclonal antibodies of the invention may be made using the hybridoma method first described by Kohler & Milstein, Nature 256:495 (1975), or may be made by recombinant DNA methods (Cabilly et al., U.S. Pat. No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as hamster, is immunized as hereinabove described to elicit lymphocytes that produce, or are capable of producing, antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized in vitro. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, pp.59-103 [Academic Press, 1986]).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 [1984]; and Brodeur et al., Monoclonal Antibody Production Techniques and Applications, pp.51-63, Marcel Dekker, Inc., New York, 1987). See, also, Boerner et al., J. Immunol., 147(1):86-95 (1991) and WO 91/17769, published Nov 28, 1991, for techniques for the production of human monoclonal antibodies.

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against Htk ligand. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an in vitro

binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson & Pollard, Anal. Biochem. 107:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods. Goding, Monoclonal Antibodies: Principles and Practice, pp.59-104 (Academic Press, 1986). Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium or RPMI-1640 medium. In addition, the hybridoma cells may be grown in vivo as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography. Alternatively, it is now possible to produce transgenic animals (e.g. mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy chain joining region (J<sub>H</sub>) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g. Jakobovits et al., Proc. Natl. Acad. Sci. USA 90:2551-255 (1993); and Jakobovits et al., Nature 362:255-258 (1993).

In a further embodiment, antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty et al., Nature, 348:552-554 (1990), using the Htk ligand (or a fragment thereof) to select for a suitable antibody or antibody fragment. Clackson et al., Nature, 352:624-628 (1991) and Marks et al., J. Mol. Biol., 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Mark et al., Bio/Technol. 10:779-783 [1992]), as well as combinatorial infection and in vivo recombination as a strategy for constructing very large phage libraries (Waterhouse et al., Nuc. Acids Res., 21:2265-2266 [1993]). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of "monoclonal" antibodies (especially human antibodies) which are encompassed by the present invention.

DNA encoding the monoclonal antibodies of the invention is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma

cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences, Morrison et al., Proc. Nat. Acad. Sci. 81, 6851 (1984), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. In that manner, "chimeric" or "hybrid" antibodies are prepared that have the binding specificity of an anti-Htk ligand monoclonal antibody herein.

Typically, such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody of the invention, or they are substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an Htk ligand and another antigen-combining site having specificity for a different antigen.

Chimeric or hybrid antibodies also may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

For diagnostic applications, the antibodies of the invention typically will be labeled with a detectable moiety. The detectable moiety can be any one which is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or  $^{125}\text{I}$ ; a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase.

Any method known in the art for separately conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., Nature 144:945 (1962); David et al., Biochemistry 13:1014 (1974); Pain et al., J. Immunol. Meth. 40:219 (1981); and Nygren, J. Histochem. and Cytochem. 30:407 (1982).

The antibodies of the present invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, Monoclonal Antibodies: A Manual of Techniques, pp.147-158 (CRC Press, Inc., 1987).

Competitive binding assays rely on the ability of a labeled standard (which may be an Htk ligand, or an immunologically reactive portion thereof) to compete with the test sample analyte (Htk ligand) for binding with a limited amount of antibody. The amount of Htk ligand in the test

sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies generally are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three part complex. See, e.g., US Pat No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

### C. Humanized Antibodies

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones et al., Nature 321:522-525 [1986]; Riechmann et al., Nature 332:323-327 [1988]; Verhoeyen et al., Science 239:1534-1536 [1988]), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (Cabilly, *supra*), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues, and possibly some FR residues, are substituted by residues from analogous sites in rodent antibodies.

It is important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three dimensional models of the parental and humanized sequences. Three dimensional immunoglobulin models are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined

from the consensus and import sequence so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding. For further details see WO 92/22653, published Dec 23, 1992.

D. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for a Htk ligand, the other one is for any other antigen, and preferably for a receptor or receptor subunit. For example, bispecific antibodies specifically binding a Htk receptor and Htk ligand are within the scope of the present invention.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Millstein and Cuello, Nature 305:537-539 [1983]). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker et al., EMBO 10:3655-3659 (1991).

According to a different and more preferred approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2 and CH3 regions. It is preferred to have the first heavy chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are cotransfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance. In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was

found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation.

5 This approach is disclosed in WO 94/04690 published March 3, 1994. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology 121:210 (1986).

#### B. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

10  
15

#### 5. Uses of Htk ligand Antibodies

Htk antibodies may be useful in certain therapeutic indications to block activity of the Htk ligand (for example in mammary carcinogenesis).

20

Therapeutic Htk ligand antibody formulations and modes for administration will be similar to those described above for Htk ligand. A typical daily dosage of the antibody might range from about 1  $\mu$ g/kg to up to 5 mg/kg or more, depending on the factors mentioned above for Htk ligand administration.

25

Htk ligand antibodies may also be useful in diagnostic assays for Htk ligand, e.g., detecting its expression in specific cells, tissues, or serum. The antibodies are labeled in the same fashion as Htk ligand described above and/or are immobilized on an insoluble matrix. Htk ligand antibodies also are useful for the affinity purification of Htk ligand from recombinant cell culture or natural sources. Htk ligand antibodies that do not detectably cross-react with other proteins can be used to purify Htk ligand free from these other known proteins. Suitable diagnostic assays for Htk ligand and its antibodies are described above.

30

#### 35 III. Experimental

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

40 All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

EXAMPLE 1Production of a Soluble Htk Receptor-Fc Fusion Protein  
for the Identification of an Htk ligand

In order to identify and ultimately clone the Htk ligand, a fusion  
5 protein is constructed which consists of the extracellular domain (ECD) of  
the Htk receptor fused to human IgG<sub>1</sub> Fc. See Bennett et al., *supra* for  
techniques for the production of the fusion protein.

The Htk receptor-Fc fusion is used to screen a series of kidney cell  
lines for their capacity to bind the extracellular domain of the Htk  
10 receptor, using FACS analysis, as previously described. See Urdal et al.,  
J. Biol. Chem. 263:2870-2877 (1988); and Gearing et al., EMBO J.  
8:3667-3676 (1989). Any cell line specifically binding the fusion protein  
is indicative of a membrane bound or membrane associated source of the Htk  
ligand. Screening of some 15 kidney cell lines results in the discovery  
15 of specific binding to one murine kidney mesangial cell line termed SV40MES  
13. The SV40MES 13 cell line is demonstrated to be positive for Htk-Fc  
binding and not for other Fc fusion proteins.

Binding competition studies are performed as follows. SV40MES 13  
cells ( $5 \times 10^6$  cells per well) are assayed for steady-state binding of  
20  $^{125}\text{I}$ -Htk-Fc in the presence of varying amounts of unlabelled Htk-Fc. The  
cells are incubated with 1 nM or 0.2 nM  $^{125}\text{I}$ -Htk-Fc and various  
concentrations of unlabelled Htk-Fc (10 pM - 1  $\mu\text{M}$ ) for 2 hr. at 4°C. Cells  
and unbound  $^{125}\text{I}$ -labelled Htk-Fc are separated by centrifugation through a  
sucrose cushion as previously described in Lee et al., J. Biol. Chem.  
25 267:16283-16287 (1992). The binding data are analyzed to determine the  
affinity and number of sites per cell as described in Munson and Rodbard  
Anal. Biochem 107:220-239 (1980). Htk-Fc fusion protein is iodinated by  
the lactoperoxidase method as described in Urdal et al., J. Biol. Chem.  
263:2870-2877 (1988). The  $K_d$  for fusion protein binding to SV40MES 13 is  
30 3 nM with approximately 6,500 sites per cell (Figure 5A). Conditioned  
media from the SV40MES 13 cell line is unable to activate tyrosine  
autophosphorylation of the Htk receptor, supporting the concept of a  
membrane bound ligand.

EXAMPLE 2Cloning the Murine Htk Ligand

The Htk receptor-Fc protein is used to expression clone the Htk  
ligand from an SV40MES 13 cDNA library transiently transfected into COS-7  
cells, as follows. A cDNA expression library from the SV40MES 13 cell line  
is constructed in the plasmid vector pRK5B (Holmes et al., Science  
40 253:1278-1280 [1991]). Fifty pools of approximately 2000 cDNAs each are  
initially transfected into COS-7 cells and the cells are screened for the  
capacity to bind Htk receptor-Fc, using slide autoradiography, as described  
in Gearing et al., EMBO J. 8:3667-3676 (1989). Five positive pools result  
from this initial screen and two of these pools are gradually subdivided  
45 in successive rounds of screening until individual clones are obtained.

Binding competition experiments are performed using one of the positive clones (#7), termed murine pRK5B-Htk ligand. In particular, binding competition curves are generated as described above with respect to SV40MES 13, using monolayers of COS-7 cells ( $5 \times 10^5$  cells per well) which are transiently transfected with clone #7, using the DEAE Dextran transfection method (McMahan et al., *EMBO J.* 10:2821-2830 [1991]).

Transfected COS-7 cells (COS-7t) used at  $2.5 \times 10^4$  cells per binding point are assayed for steady-state binding of  $^{125}\text{I}$ -Htk-Fc in the presence of varying amounts of unlabelled Htk-Fc, as described above. Htk-Fc binding to transfected COS-7 cells demonstrates a  $K_d$  of 500 pm (Figure 5B), indicating that clone #7 is the murine Htk-ligand.

The DNA sequence and deduced amino acid sequence of the murine Htk ligand are shown in Figures 1A-B. The predicted molecular weight of the protein following signal peptide cleavage is 34 kD with an estimated pI of 8.9.

The sequence derived from the #7 clone is confirmed by sequencing another independent clone of 4700 bp that gives the identical coding sequence. DNA sequencing is performed using the ABI Taq Dye Deoxy terminator cycle sequencing kit on an automated Applied Biosystems DNA sequencer, model 373A. Both strands of individual clones are sequenced in their entirety.

Sequence comparison of the Htk ligand and B61 (Bartley et al., *supra* and Holzman et al., *supra*) indicates 23% similarity between the molecules. However, B61 does not contain a transmembrane domain. Nevertheless, the degree of homology suggests that the Htk ligand and B61 may comprise members of a structurally similar family that bind to various members of the Eph/ELK family of receptor tyrosine kinases.

### EXAMPLE 3

#### Tissue Distribution of the Htk ligand

Northern blot analysis is performed in order to detect the presence of the Htk ligand in mouse adult tissues, human adult tissues and human fetal tissues. In particular, Northern blots are obtained from Clontech (Palo Alto, CA) which contain 2  $\mu\text{g}$ /lane of polyA-selected RNA from mouse adult, human adult and human fetal tissues. Mouse blots are hybridized in 50% formamide at 42°C to  $^{32}\text{P}$  labelled murine Htk ligand cDNA and washed under stringent conditions (final wash: 0.2 x SSC, 0.2% SDS at 60°C). Human tissue blots are hybridized in 35% formamide at 42°C and washed under stringent conditions as above.

Northern analysis of mouse and human Htk ligand messenger RNA in adult and fetal tissues shows only one transcript at approximately 5.2 kb which displays widespread tissue expression. In particular, the Htk ligand is present in large amounts in mouse adult lung, brain, heart and kidney, and in lesser amounts in spleen, liver, skeletal muscle and testis. The ligand is present in human adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary,



small intestine and colon but is not detectable by Northern analysis in peripheral blood leukocytes. Finally, the transcript is detectable in highest amounts in human fetal brain, lung, and kidney tissues with lower amounts detectable in human fetal liver tissue.

5        *In situ* hybridizations are also performed to detect Htk ligand DNA expression. Mouse embryos (embryonic day 13) or brains from postnatal day one or adult mice are prepared for *in situ* hybridization as follows. Freshly dissected brains or embryos fixed in 4% formaldehyde are frozen and sectioned with a cryostat. Sections are thaw-mounted onto slides, air  
10        dried, and stored at -70°C. Hybridization is conducted with <sup>32</sup>P-labelled riboprobes by a modification of published methods (Phillips et al., Science 250:290-294 [1990]). Sense (control) and antisense cRNA probes corresponding to nucleotides 1597 to 2198 of the murine Htk ligand DNA sequence of Figures 1A-B are utilized for hybridization.

15        On the day of hybridization, sections are brought to room temperature, fixed for 10-30 minutes in 4% formaldehyde with or without the addition of 1% glutaraldehyde in 0.1M phosphate (pH 7.2), rinsed, and incubated in hybridization buffer for 1-3 hours at 42°C. Hybridization  
20        buffer consists of 50% formamide, 0.1M NaCl, 20mM Tris HCl, pH 8.0, 1X Denhardt's solution, 10% Dextran sulfate, 10mM DTT. Probes are heated to 95°C for 3 minutes in the presence of carrier RNA, after which they are immediately cooled to 4°C. Probe is then added to the hybridization buffer on each slide at a final concentration of 6.5 X 10<sup>6</sup> cpm/ml, and allowed to hybridize at 55°C overnight. Following hybridization, sections are treated  
25        as follows: 2 rinses in 2X SSC, 30 minute incubation in RNase A (20 µg/ml), 2 rinses in 2X SSC, 1 hour incubation at 55°C in 0.1X SSC, 2 rinses in 0.5X SSC, dehydration in a series of graded ethanol solutions (60%, 75%, and 85% ethanol containing 0.3 M ammonium acetate, followed by 90% and 100% ethanol), and air drying at room temperature. The sections are then  
30        exposed to sheet film (Beta-Max, Amersham) for a period of 1 to 3 days after which they are dipped in emulsion (Amersham LM-1) and exposed at 4°C for 3 to 8 weeks. Film and emulsion autoradiographs are developed by treatment with standard photographic developer and fixer.

35        The sheet film autoradiographs are viewed both by visual inspection on a light box and through a stereoscopic microscope. The emulsion autoradiographs are viewed under both brightfield and darkfield microscopy. Observation of the autoradiograms reveals hybridization signal in several regions on the sections hybridized with antisense probe that were not observed on the control sections hybridized with sense probe. These  
40        regions include, but are not restricted to, several regions of the adult forebrain, including the CA1 region of the hippocampus, the cerebral cortex (including piriform and entorhinal cortices), and the caudate-putamen. Prominent hybridization is also observed in the cerebellar cortex. Hybridization is less intense or absent from other brain structures  
45        including the septum, white matter tracts such as the corpus callosum, and numerous diencephalic, mesencephalic, and myelencephalic regions. In the

embryo, strong hybridization is seen in (but not confined to) the developing lung, digestive tract, liver, kidney, salivary gland, vertebrae, muscle, olfactory epithelium, epithelium of developing ear, within both dorsal root and trigeminal ganglia, meninges of both brain and spinal cord, and within numerous regions of both brain and spinal cord. Within the developing brain, expression is notably intense in the developing forebrain, but significant hybridization was observed in all major subdivisions (telencephalon, diencephalon, mesencephalon, metencephalon, and myelencephalon).

10

EXAMPLE 4Induction of Tyrosine Phosphorylation of the  
Htk Receptor by the Htk Ligand

To determine whether the Htk ligand stimulates Htk phosphorylation, and to further confirm that clone #7 described above indeed codes for a ligand for the Htk receptor, the following experiment is performed.

NIH 3T3 cells are stably transfected with the full-length Htk receptor. A 4038 bp ClaI - XbaI cDNA fragment containing 32 bp of linker sequence, 37 bp of pBluescript (Stratagene La Jolla, CA) polylinker and the entire 3969 bp Htk receptor cDNA is subcloned into the expression vector pRIS (Genentech, Inc.) under the control of the Rous sarcoma virus LTR promoter. NIH3T3 cells maintained in high glucose Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FCS are co-transfected with pRIS-Htk receptor and pNeo (an SV40 based vector containing the neomycin resistance marker) by the calcium phosphate method as described by Gorman et al., in DNA Prot. Engineer. Tech. 2:3-10 [1990]. Neomycin resistant colonies are selected 48 hours after transfection with Geneticin (Gibco/BRL) at 400 µg/ml. Fourteen days later individual resistant colonies are isolated, expanded and analyzed by flow cytometry for Htk receptor expression using rabbit polyclonal antiserum. Specificity of response is demonstrated using mock-transfected 3T3 cells.

Then, one million transfected 3T3 cells (3T3-T) or non-transfected 3T3 cells (3T3) are co-incubated with  $1 \times 10^6$  Htk ligand transiently transfected COS-7 cells (transfected with clone #7 from above using the DEAE-dextran method as described above), mock-transfected COS-7 cells, or  $3 \times 10^6$  SV40MES 13 cells, at 37°C for 30 minutes. Transfected and mock transfected NIH 3T3 cells are also incubated with monoclonal anti-human Htk receptor antibody (IC2-C2) produced by hybridoma Anti-HpTK 5 (ATCC Accession No. HB 11,583), known to induce autophosphorylation of the Htk receptor.

Cells are lysed in NP-40 lysis buffer (1% NP-40, 1 mM EDTA, 200 mM NaCl, 50 mM Tris Cl, pH 8.0, 2 mM DMSF, 2.5 mM  $\text{Na}_2\text{VO}_4$ ) and immunoprecipitated with anti-human Htk rabbit polyclonal sera, produced as follows. Polyclonal antibodies are generated in New Zealand White rabbits against the soluble Htk receptor-Fc fusion protein described in Bennett et al., supra. 4 µg of the protein in 100 µl PBS is emulsified with 100 µl

Freund's adjuvant (complete adjuvant for the primary injection and incomplete adjuvant for all boosts). For the primary immunization and the first boost, the protein is injected directly into the popliteal lymph nodes (Sigel et al., Methods Enzymol. 93:3-12 [1983]). For subsequent  
 5 boosts, the protein is injected into subcutaneous and intramuscular sites. 1.3 µg protein/kg body weight is injected every 3 weeks with bleeds taken 1 and 2 weeks following each boost. Specificity of the antibody is demonstrated by flow cytometric analysis of NIH3T3 cells transfected with full length Htk receptor or vector alone using a 1:200 dilution of the pre-  
 10 immune serum or anti-Htk receptor-IgG Fc serum.

Immunoprecipitated cells are analyzed on SDS-PAGE 4-12% gradient gels. Gels are then transferred to nitrocellulose filters and Western blotted using the antiphosphotyrosine antibody 4G10 (UBI, Lake Placid, New York). Both clone #7 transfected COS-7 cells, SV40MES 13 cells, and IC2-C2  
 15 antibody induce autophosphorylation of the Htk receptor upon coincubation, confirming that the Htk ligand stimulates Htk phosphorylation and that clone #7 codes for a Htk ligand.

#### EXAMPLE 5

##### Cloning the Human Htk ligand

20 In order to clone the human Htk ligand, a human fetal brain cDNA library is prepared using the techniques generally described in Sambrook et al., *supra*. A human fetal lung library is purchased from Clontech (Palo Alto, CA). These libraries are screened using techniques described in Sambrook et al., *supra*, with a fragment from the 5' end of the mouse cDNA  
 25 as a probe (*i.e.*, residues 515 to 2,312 of Figures 1A-B). The entire human Htk ligand gene is found to be present in a single clone isolated from the human fetal brain library. The plasmid having the nucleic acid encoding the human Htk ligand has been deposited with the American Type Culture Collection (ATCC) on June 24, 1994 under Accession No. 75,820. The  
 30 nucleotide and amino acid sequences of the human Htk ligand are shown in Figure 2. The sequence encodes a protein having a predicted molecular weight of 34kD following signal peptide cleavage. The murine and human ligands show 96% sequence identity at the amino acid level, demonstrating a high degree of conservation between species. This is consistent with the  
 35 homology between human Htk receptor and its mouse homologue, myk-1, which are 91% identical at the amino acid level.

#### Deposits

The following cultures have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, USA (ATCC):

40	<u>Hybridoma</u>	<u>ATCC No.</u>	<u>Deposit Date</u>
	Anti-HpTK5	HB 11,583	March 15, 1994
	Plasmid Human	ATCC 75,820	June 24, 1994
	Htk Ligand		

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of viable cultures for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the cultures to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the cultures to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if the cultures should die or be lost or destroyed when cultivated under suitable conditions, they will be promptly replaced on notification with a viable specimen of the same culture. Availability of the deposited strains is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the cultures deposited, since the deposited embodiment is intended as a single illustration of one aspect of the invention and any cultures that are functionally equivalent are within the scope of this invention. The deposit of materials herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustration that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Genentech, Inc.
- (ii) TITLE OF INVENTION: HTK LIGAND
- 10 (iii) NUMBER OF SEQUENCES: 7
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Genentech, Inc.
- (B) STREET: 460 Point San Bruno Blvd
- 15 (C) CITY: South San Francisco
- (D) STATE: California
- (E) COUNTRY: USA
- (F) ZIP: 94080
- 20 (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: 5.25 inch, 360 Kb floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: patin (Genentech)
- 25 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (C) CLASSIFICATION:
- 30 (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (viii) ATTORNEY/AGENT INFORMATION:
- 35 (A) NAME: Lee, Wendy M.
- (B) REGISTRATION NUMBER: 00,000
- (C) REFERENCE/DOCKET NUMBER: 902PCT
- (ix) TELECOMMUNICATION INFORMATION:
- 40 (A) TELEPHONE: 415/225-1994
- (B) TELEFAX: 415/952-9881
- (C) TELEX: 910/371-7168

## (2) INFORMATION FOR SEQ ID NO:1:

- 45 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 4342 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 50 (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
- 55 CCCACGCGTC CGCGCGCGCT GAGGGACGCG CAGGGTGAGC GCACCTGGCC 50
- TGGGCGACCG CGGGAGCGGC GCGGCGTGTC CGCCCCGGAG GATTGGGGGT 100
- 60 CGCTGCCCCGCG GGCCGGTCCC AACGCGTCCC GGAGTGCGCA GAACTGGGAG 150
- CGGCTTGGGC ATGGCCATGG CCCGGTCCAG GAGGGACTCT GTGTGGAAGT 200

ACTGTTGGGG ACTTTTGATG GTTTTGTGCA GAACTGCGAT CTCCAGATCG 250

5 ATAGTTTTAG AGCCTATCTA CTGGAATTCC TCGAACTCCA AATTTCTACC 300

CGGACAAGGC CTGGTACTAT ACCCACAGAT AGGAGACAAA TTGGATATTA 350

10 TTTGCCCCAA AGTGGACTCT AAAACTGTTG GCCAGTATGA ATATTATAAA 400

15 GTTTATATGG TTGATAAAGA CCAAGCAGAC AGATGCACAA TTAAGAAGGA 450

GAATACCCCG CTGCTCAACT GTGCCAGACC AGACCAAGAT GTGAAATTCA 500

20 CCATCAAGTT TCAAGAATTC AGCCCTAACC TCTGGGGTCT AGAATTTAG 550

AAGAACAAAG ATTACTACAT TATATCTACA TCAAATGGGT CTTTGGAGGG 600

25 CCTGGATAAC CAGGAGGGAG GGGTGTGCCA GACAAGAGCC ATGAAGATCC 650

30 TCATGAAAGT TGGACAAGAT GCAAGTTCTG CTGGATCAGC CAGGAATCAC 700

GGTCCAACAA GACGTCCAGA GCTAGAAGCT GGTACAAATG GGAGAAGTTC 750

35 AACAACAAGT CCCTTTGTGA AGCCAAATCC AGGTTCTAGC ACCGATGGCA 800

ACAGCGCGGG GCATTCCGGG AACAACTCTC TGGGTTCGGA AGTGGCCTTA 850

40 TTCGCAGGGA TCGCATCAGG ATGCATCATC TTCATCGTCA TCATCATCAC 900

TTTGGTGGTG CTGCTGCTCA AGTACCGCAG GAGACACCGC AAACACTCTC 950

45 CACAGCACAC GACCACGCTG TCTCTCAGCA CACTGGCCAC GCCCAAGCGA 1000

50 GGTGGCAACA ACAATGGCTC GGAGCCAGT GACGTTATCA TACCACTAAG 1050

GACTGCAGAC AGCGTCTTCT GCCCGCACTA CGAGAAGGTC AGCGGGGACT 1100

55 ATGGGCACCC GGTGTACATC GTGCAGGAGA TGCCCCACA GAGTCCTGCC 1150

AACATTTACT ACAAGGTCTG AGGCCTGAGA CCTGCGCCTC CCAAGGGAAC 1200

60 TCGCACCTTG TTCTTGGGCA CGCAGGGACT GCCTGAGCCT GGCTGTGGGG 1250

GCAGGATGCC TCCTGGAAGA GCCTGGATCT GGACAGTTTT GTAGTCTGTA 1300

5 GCTTTTCCGA CCCTGGGGAC CACAGACCCT CCCCGBAAGC TGGAAGACTG 1350

CTAGGAGATC CCCACTTGA CTGCCGCGGC CCCACGCGGA CCTCCAAGCC 1400

10 ATGCACCCAG CCACTCAGGC CTCTGCAGAG CCCGGGGAGG ACACGGTAGG 1450

CTATGGATGG CGCAGCAGCA TCTTAGGAGA AGGTTGCGCA CCAGGCCGGC 1500

15 CCCTGCCTCC ACGTTTCCTG CCGTGCACAC TGGACTTATC ACTTGGACCT 1550

CGGGTTCAGT AAGGTTTTCA AAGATCTCTA GTGTTTAGTC CTCACTCACT 1600

20 CACTCACTCA CTCACTCCTT CTCTTGCCAG GGCTCTGCAG CAAACTCCCT 1650

AGACCCCTCA CTCCACGTAC TGCATCATTA CGGGACACTC ACCACAGAGT 1700

CCCAGCTCCA CCCTTTACAC CAAGATCAAA TTAGATGGGT ATTAGGTACA 1750

30 GAAGAACCCT GCTTGCCTGG AGGCCGGGTC AGCCGGGAAG CGCAGATGTG 1800

TGGAGGAGTG AGGAGGTGCT GGCTGCCACG GCAGGTCAAG GCTGCTTGCT 1850

35 GCCCCTGGAG CATAGTAGGG ATGCAGGAAG GAAATAGATA ATGCTTTGGT 1900

TTTTCTGAGA GGACAGAGAC AGGTGGGAGG TGAAGTACTG GTGAGTGGTG 1950

GGGAGCCTTT CACTACCACA CAGCTATGCA GCAGGGAATC AAAAGTCCCT 2000

45 CTCCTGCGGG GAACAAAGGG GCCATTGTTG TGAAAGGACC AGCTAGAGCA 2050

CAGAGGGAGA GGGCAGGCCT CCGGTGAAGT GCTGGGGCAG AACTGCAGAG 2100

50 GTACTGGAAA TAAAAAGCGC AGCGCAGAGC TGTGGGAGAG TCCGTCTGCT 2150

TTGGGAGATG TTTTAAGCAG ACTCAGCTGC TATATTACCA CGTTTTTATT 2200

AAAAACACAG GGAAAGCATT TAGGAGAAGA GCAGAGAGCC AAATCTGACC 2250

60 TAGAAGTTGA AAAGCCAAAG GTCAAACAGG CTGTAAGTCC ATCACCCTG 2300

AGGTTATTGG AGAATTCTCA TTAGGAAGG CAGGTCAGAT TCCCCAGGCC 2350

CCATAAGTGC CCCTTCCCCC TCCCTGATTG AGCCTTACAC GTTGGTTTTT 2400

5 GGTTTATGGC CGTGCTGTCC GGGCTCCAAG GCAGTACCCG GGCTCCATGT 2450

CAAAGCAAAG CACACATGGC CCACCTCTTA GAGTCCTTGA GATGGAAGTA 2500

10 AGTTATGCCG CGGAAGGAAA GGCGAAGATA GGACATATTT ATAATAGGTG 2550

TATAGAACAC AAGGGATATA AAATGAAAGA TTTTACTAA TATATATTTT 2600

15 AAGATTACAC ACAATACACA CCAGAAGACG TGGAGTTCGG TGGTGGTGGT 2650

20 GGTCGTGGTG GTGGTGATTA AAGTGACCCC AGCGCTTAGT GCTTTAAAAA 2700

GTGAAAGATT GGGTAGCTAC TCCCCGAAAC GTACCAATAG CAAGAAAAGT 2750

25 ATCCATAATG AGAGCAAATG GCAAAAATAA CACGGTCCTG CGGGAATCTC 2800

GCAGAACCGT AGACTAGGAA TGCCAGCCCC CCAAATTGAT GTGACCCTGC 2850

30 CCCGGGTTAG ACAATGATAA AATGCGCTGG CCTTTATTTT CTGTGTTGGG 2900

35 TTTTCCCTTG CCTTATGGGC TGAAGTGTTT TCTAGAATTT AGCAGGTCAC 2950

ACTGAGGGGA TTCCAGTTTA ACTGTGGGTC CCTCCTCCTC TCCTACCCCA 3000

40 TCCCTGCCCT TCCAGAGAAT AACAGGAAGC CTTCTTTTTT TTTTTTTTTT 3050

AAGTGCTATG CAAAAGAGAC ATCTTTAACA GAGTCCTGTT ACTATGGTAA 3100

45 CATTTTGCTT TCTGAATTGG GAGGAAATAA AAATTGTAAT GACAGCATTT 3150

50 GAAGGTTCTC AGACCTCCAG TGAGTACCTG CAAAAATGAG TTGTCACAGA 3200

GATTATTCCC TACTTCTCAA ACCTGAAAAT GATGTTGGTT CGATGTGTGT 3250

55 GTGTGTGTAT GAGTGGGTGT GTGGTACATG TGTGTACATA TATGTATAAT 3300

ATATATCTCC AGTATATATT ATATATATCT ATATCATATT TCTGTGGAGG 3350

60 GTTGCCATGG CAATCAACTG CAGTACATAT GTAGTTCTTT CCATCACCCCT 3400



5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

AACCTCTCCT GCGCATTAC ACAAGAGTTT CTTGTAAGCC ATCAAAAGTT 3450  
AATTCTAGGG GGAGAGGGAT GAGGCGGGGA GACATGGGAA ACCGTCTGAT 3500  
TTTAATGAAA TCAAATGTCT GTGTCATCGG TTGGCTACGT TTTGGTTCTA 3550  
TGCTAAACTG TGAAGAATCG GATGAATTGA TGAAGAGTTG AGTTACCTGC 3600  
AACCATTGA GAAGTGTCTT GTGCGTCTGT TTTGTGTCTG GTGCAGAAAA 3650  
TGACAATCTA CCAACTGTCC CCTTATTTGG AGTTGGTTCA GCTTTGGAAA 3700  
GTTACTGTAA ATGCCTTGCT TGTATTATCA TCCCTAGTCA CCTGACTTCG 3750  
GAGCTTGAC CATCGTGTTC TAAGTGAAGA CGCTGTAAAT AGGTTTCAGAT 3800  
CTTACCGTCT ATGGATTCGG GTGTTACAGT AGCCTTATTC ACCTTTTTTAA 3850  
TAAAAATACA CATGAAAACG AGACAGTAAT GGCTTTTCTT ACCCAGATTG 3900  
TGTACATAGA GCAATGTTGG TTTTTTATAA AGTCTAAGCA AGATGTTTTG 3950  
TATAAATCT GAATTTTGCA ATGTATTTAG CTACAGCTTT TAACGGCAGT 4000  
GTCATCCCCT TTGCACTGTA ATGAGGAAAA AAAAAAGGTA TAAAAGGTTG 4050  
CCAAATTGCT GCATATTTGT GCCGTAATTA TGTACCATGA ATATTTATTT 4100  
AATTTTCGTTG TCCAATTTGT AAGTAACACA GTATTATGCT TGAGTTATAA 4150  
ATATTTTTTC TTTGTTTTAT TTTAATAGCC TGTCATAGGT TTTTTTTTAA 4200  
TCTGCTTTAG TTCCACATGA CAGTTAAGCC CCAGAAATGA GATCCGAGCA 4250  
GCCACATTCC ACGTCTGTTT CAAAATGAAT TTGTTCTTAA AAAAAATAAA 4300  
ATATTTTTTTT CCTATGGAAA AAAAAAAAAA AAGGGCGGCC GC 4342

## (2) INFORMATION FOR SEQ ID NO:2:

60

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 336 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	Met	Ala	Met	Ala	Arg	Ser	Arg	Arg	Asp	Ser	Val	Trp	Lys	Tyr	Cys	
	1				5					10					15	
5	Trp	Gly	Leu	Leu	Met	Val	Leu	Cys	Arg	Thr	Ala	Ile	Ser	Arg	Ser	
					20					25					30	
10	Ile	Val	Leu	Glu	Pro	Ile	Tyr	Trp	Asn	Ser	Ser	Asn	Ser	Lys	Phe	
					35					40					45	
	Leu	Pro	Gly	Gln	Gly	Leu	Val	Leu	Tyr	Pro	Gln	Ile	Gly	Asp	Lys	
					50					55					60	
15	Leu	Asp	Ile	Ile	Cys	Pro	Lys	Val	Asp	Ser	Lys	Thr	Val	Gly	Gln	
					65					70					75	
	Tyr	Glu	Tyr	Tyr	Lys	Val	Tyr	Met	Val	Asp	Lys	Asp	Gln	Ala	Asp	
					80					85					90	
20	Arg	Cys	Thr	Ile	Lys	Lys	Glu	Asn	Thr	Pro	Leu	Leu	Asn	Cys	Ala	
					95					100					105	
25	Arg	Pro	Asp	Gln	Asp	Val	Lys	Phe	Thr	Ile	Lys	Phe	Gln	Glu	Phe	
					110					115					120	
	Ser	Pro	Asn	Leu	Trp	Gly	Leu	Glu	Phe	Gln	Lys	Asn	Lys	Asp	Tyr	
					125					130					135	
30	Tyr	Ile	Ile	Ser	Thr	Ser	Asn	Gly	Ser	Leu	Glu	Gly	Leu	Asp	Asn	
					140					145					150	
	Gln	Glu	Gly	Gly	Val	Cys	Gln	Thr	Arg	Ala	Met	Lys	Ile	Leu	Met	
					155					160					165	
35	Lys	Val	Gly	Gln	Asp	Ala	Ser	Ser	Ala	Gly	Ser	Ala	Arg	Asn	His	
					170					175					180	
40	Gly	Pro	Thr	Arg	Arg	Pro	Glu	Leu	Glu	Ala	Gly	Thr	Asn	Gly	Arg	
					185					190					195	
	Ser	Ser	Thr	Thr	Ser	Pro	Phe	Val	Lys	Pro	Asn	Pro	Gly	Ser	Ser	
					200					205					210	
45	Thr	Asp	Gly	Asn	Ser	Ala	Gly	His	Ser	Gly	Asn	Asn	Leu	Leu	Gly	
					215					220					225	
	Ser	Glu	Val	Ala	Leu	Phe	Ala	Gly	Ile	Ala	Ser	Gly	Cys	Ile	Ile	
					230					235					240	
50	Phe	Ile	Val	Ile	Ile	Ile	Thr	Leu	Val	Val	Leu	Leu	Leu	Lys	Tyr	
					245					250					255	
	Arg	Arg	Arg	His	Arg	Lys	His	Ser	Pro	Gln	His	Thr	Thr	Thr	Leu	
					260					265					270	
55	Ser	Leu	Ser	Thr	Leu	Ala	Thr	Pro	Lys	Arg	Gly	Gly	Asn	Asn	Asn	
					275					280					285	
60	Gly	Ser	Glu	Pro	Ser	Asp	Val	Ile	Ile	Pro	Leu	Arg	Thr	Ala	Asp	
					290					295					300	
	Ser	Val	Phe	Cys	Pro	His	Tyr	Glu	Lys	Val	Ser	Gly	Asp	Tyr	Gly	
					305					310					315	

His Pro Val Tyr Ile Val Gln Glu Met Pro Pro Gln Ser Pro Ala  
320 325 330

5 Asn Ile Tyr Tyr Lys Val  
335 336

(2) INFORMATION FOR SEQ ID NO:3:

- 10 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 1953 bases  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTCAGGCGC GGAGCTGGGA GTGGCTTCGC CATGGCTGTG AGAAGGGACT 50  
20 CCGTGTGGAA GTACTGCTGG GGTGTTTTGA TGGTTTTATG CAGAACTGCG 100  
ATTTCCAAAT CGATAGTTTT AGAGCCTATC TATTGGAATT CCTCGAACTC 150  
25 CAAATTTCTA CCTGGACAAG GACTGGTACT ATACCCACAG ATAGGAGACA 200  
AATTGGATAT TATTTGCCCC AAAGTGGACT CTAAACTGT TGGCCAGTAT 250  
GAATATTATA AAGTTTATAT GGTTGATAAA GACCAAGCAG ACAGATGCAC 300  
35 TATTAAGAAG GAAAATACCC CTCTCCTCAA CTGTGCCAAA CCAGACCAAG 350  
ATATCAAATT CACCATCAAG TTTCAAGAAT TCAGCCCTAA CCTCTGGGGT 400  
40 CTAGAATTC AGAAGAACAA AGATTATTAC ATTATATCTA CATCAAATGG 450  
GTCTTTGGAG GGCCTGGATA ACCAGGAGGG AGGGGTGTGC CAGACAAGAG 500  
CCATGAAGAT CCTCATGAAA GTTGGACAAG ATGCAAGTTC TGCTGGATCA 550  
50 ACCAGGAATA AAGATCCAAC AAGACGTCCA GAACTAGAAG CTGGTACAAA 600  
TGGAAGAAGT TCGACAACAA GTCCCTTTGT AAAACCAAAT CCAGGTTCTA 650  
55 GCACAGACGG CAACAGCGCC GGACATTCGG GGAACAACAT CCTCGGTTCC 700  
GAAGTGGCCT TATTTGCAGG GATTGCTTCA GGATGCATCA TCTTCATCGT 750  
CATCATCATC ACGCTGGTGG TCCTCTTGCT GAAGTACCGG AGGAGACACA 800

GGAAGCACTC GCCGCAGCAC ACGACCACGC TGTCGCTCAG CACACTGGCC 850

ACACCCAAGC GCAGCGGCAA CAACAACGGC TCAGAGCCCA GTGACATTAT 900

5 CATCCCGCTA AGGACTGCGG ACAGCGTCTT CTGCCCTCAC TACGAGAAGG 950

TCAGCGGGGA CTACGGGCAC CCGGTGTACA TCGTCCAGGA GATGCCCCCG 1000

10 CAGAGCCCCG CGAACATTTA CTACAAGGTC TGAGAGGGAC CCTGGTGGTA 1050

15 CCTGTGCTTT CCCAGAGGAC ACCTAATGTC CCGATGCCTC CCTTGAGGGT 1100

TTGAGAGCCC GCGTGCTGGA GAATTGACTG AAGCACAGCA CCGGGGGAGA 1150

20 GGGACACTCC TCCTCGGAAG AGCCCGTCGC GCTGGACAGC TTACCTAGTC 1200

25 TTGTAGCATT CGGCCTTGGT GAACACACAC GCTCCCTGGA AGCTGGAAGA 1250

CTGTGCAGAA GACGCCCAT TCGGACTGCTG TGCCGCGTCC CACGTCTCCT 1300

30 CCTCGAAGCC ATGTGCTGCG GTCCTCAGG CCTCTGCAGA AGCCAAGGGA 1350

AGACAGTGGT TTGTGGACGA GAGGGCTGTG AGCATCCTGG CAGGTGCCCC 1400

35 AGGATGCCAC GCCTGGAAGG GCCGGCTTCT GCCTGGGGTG CATTTCCCCC 1450

40 GCAGTGCATA CCGGACTTGT CACACGGACC TCGGGCTAGT TAAGGTGTGC 1500

AAAGATCTCT AGAGTTTAGT CCTTACTGTC TCACTCGTTC TGTTACCCAG 1550

45 GGCTCTGCAG CACCTCACCT GAGACCTCCA CTCCACATCT GCATCACTCA 1600

TGGAACACTC ATGTCTGGAG TCCCCTCCTC CAGCCGCTGG CAACAACAGC 1650

50 TTCAGTCCAT GGGTAATCCG TTCATAGAAA TTGTGTTTGC TAACAAGGTG 1700

55 CCCTTTAGCC AGATGCTAGG CTGTCTGCGA AGAAGGCTAG GAGTTCATAG 1750

AAGGGAGTGG GGCTGGGGAA AGGGCTGGCT GCAATTGCAG CTCCTGCTG 1800

60 CTGCCTCTGA AACAGAAAGT TGGAAGGAA AAAAGAAAAA AGCAATTAGG 1850

TAGCACAGCA CTTTGGTTTT GCTGAGATCG AAGAGGCCAG TAGGAGACAC 1900

GACAGCACAC ACAGTGGATT CCAGTGCATG GGGAGGCGGT CGACGAGCTC 1950

5 GAG 1953

(2) INFORMATION FOR SEQ ID NO:4:

10

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 333 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Ala	Val	Arg	Arg	Asp	Ser	Val	Trp	Lys	Tyr	Cys	Trp	Gly	Val	1	5	10	15
Leu	Met	Val	Leu	Cys	Arg	Thr	Ala	Ile	Ser	Lys	Ser	Ile	Val	Leu	20	25	30	
Glu	Pro	Ile	Tyr	Trp	Asn	Ser	Ser	Asn	Ser	Lys	Phe	Leu	Pro	Gly	25	35	40	45
Gln	Gly	Leu	Val	Leu	Tyr	Pro	Gln	Ile	Gly	Asp	Lys	Leu	Asp	Ile	50	55	60	
Ile	Cys	Pro	Lys	Val	Asp	Ser	Lys	Thr	Val	Gly	Gln	Tyr	Glu	Tyr	30	65	70	75
Tyr	Lys	Val	Tyr	Met	Val	Asp	Lys	Asp	Gln	Ala	Asp	Arg	Cys	Thr	35	80	85	90
Ile	Lys	Lys	Glu	Asn	Thr	Pro	Leu	Leu	Asn	Cys	Ala	Lys	Pro	Asp	95	100	105	
Gln	Asp	Ile	Lys	Phe	Thr	Ile	Lys	Phe	Gln	Glu	Phe	Ser	Pro	Asn	40	110	115	120
Leu	Trp	Gly	Leu	Glu	Phe	Gln	Lys	Asn	Lys	Asp	Tyr	Tyr	Ile	Ile	125	130	135	
Ser	Thr	Ser	Asn	Gly	Ser	Leu	Glu	Gly	Leu	Asp	Asn	Gln	Glu	Gly	45	140	145	150
Gly	Val	Cys	Gln	Thr	Arg	Ala	Met	Lys	Ile	Leu	Met	Lys	Val	Gly	155	160	165	
Gln	Asp	Ala	Ser	Ser	Ala	Gly	Ser	Thr	Arg	Asn	Lys	Asp	Pro	Thr	50	170	175	180
Arg	Arg	Pro	Glu	Leu	Glu	Ala	Gly	Thr	Asn	Gly	Arg	Ser	Ser	Thr	55	185	190	195
Thr	Ser	Pro	Phe	Val	Lys	Pro	Asn	Pro	Gly	Ser	Ser	Thr	Asp	Gly	200	205	210	
Asn	Ser	Ala	Gly	His	Ser	Gly	Asn	Asn	Ile	Leu	Gly	Ser	Glu	Val	60	215	220	225
Ala	Leu	Phe	Ala	Gly	Ile	Ala	Ser	Gly	Cys	Ile	Ile	Phe	Ile	Val	230	235	240	

Ile Ile Ile Thr Leu Val Val Leu Leu Leu Lys Tyr Arg Arg Arg  
 245 250 255  
 5 His Arg Lys His Ser Pro Gln His Thr Thr Thr Leu Ser Leu Ser  
 260 265 270  
 Thr Leu Ala Thr Pro Lys Arg Ser Gly Asn Asn Asn Gly Ser Glu  
 275 280 285  
 10 Pro Ser Asp Ile Ile Ile Pro Leu Arg Thr Ala Asp Ser Val Phe  
 290 295 300  
 Cys Pro His Tyr Glu Lys Val Ser Gly Asp Tyr Gly His Pro Val  
 305 310 315  
 15 Tyr Ile Val Gln Glu Met Pro Pro Gln Ser Pro Ala Asn Ile Tyr  
 320 325 330  
 Tyr Lys Val  
 20 333

## (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:  
 25 (A) LENGTH: 3969 bases  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TCGGCGTCCA CCCGCCAGG GAGAGTCAGA CCTGGGGGGG CGAGGGCCCC 50  
 35 CCAAACCTCAG TTCGGATCCT ACCCGAGTGA GCGGGCGCCA TGGAGCTCCG 100  
 GGTGCTGCTC TGCTGGGCTT CGTTGGCCGC AGCTTTGGAA GAGACCCTGC 150  
 40 TGAACACAAA ATTGGAAACT GCTGATCTGA AGTGGGTGAC ATTCCCTCAG 200  
 45 GTGGACGGGC AGTGGGAGGA ACTGAGCGGC CTGGATGAGG AACAGCACAG 250  
 CGTGCGCACC TACGAAGTGT GTGACGTGCA GCGTGCCCCG GGCCAGGCCC 300  
 50 ACTGGCTTCG CACAGGTTGG GTCCCACGGC GGGGCGCCGT CCACGTGTAC 350  
 GCCACGCTGC GCTTCACCAT GCTCGAGTGC CTGTCCCTGC CTCGGGCTGG 400  
 55 GCGCTCCTGC AAGGAGACCT TCACCGTCTT CTA CTATGAG AGCGATGCGG 450  
 60 ACACGGCCAC GGCCCTCAG CCAGCCTGGA TGGAGAACCC CTACATCAAG 500  
 GTGGACACGG TGGCCGCGGA GCATCTCACC CGGAAGCGCC CTGGGGCCGA 550

GGCCACCGGG AAGGTGAATG TCAAGACGCT GCGTCTGGGA CCGCTCAGCA 600

5 AGGCTGGCTT CTACCTGGCC TTCCAGGACC AGGGTGCCTG CATGGCCCTG 650

CTATCCCTGC ACCTCTTCTA CAAAAAGTGC GCCCAGCTGA CTGTGAACCT 700

10 GACTCGATT CCGGAGACTG TGCCTCGGGA GCTGGTTGTG CCCGTGGCCG 750

GTAGCTGCGT GGTGGATGCC GTCCCCGCCC CTGGCCCCAG CCCCAGCCTC 800

15 TACTGCCGTG AGGATGGCCA GTGGGCCGAA CAGCCGGTCA CGGGCTGCAG 850

CTGTGCTCCG GGGTTCGAGG CAGCTGAGGG GAACACCAAG TGCCGAGCCT 900

20 GTGCCCAGGG CACCTTCAAG CCCCTGTCAG GAGAAGGGTC CTGCCAGCCA 950

25 TGCCCAGCCA ATAGCCACTC TAACACCATT GGATCAGCCG TCTGCCAGTG 1000

CCGCGTCGGG TACTTCCGGG CACGCACAGA CCCCCGGGGT GCACCCTGCA 1050

30 CCACCCCTCC TTCGGCTCCG CGGAGCGTGG TTTCCCGCCT GAACGGCTCC 1100

TCCCTGCACC TGGAAATGGAG TGCCCCCCTG GAGTCTGGTG GCCGAGAGGA 1150

35 CCTCACCTAC GCCCTCCGCT GCCGGGAGTG CCGACCCGGA GGCTCCTGTG 1200

40 CGCCCTGCGG GGGAGACCTG ACTTTTGACC CCGGCCCCCG GGACCTGGTG 1250

GAGCCCTGGG TGGTGGTTCG AGGGCTACGT CCTGACTTCA CCTATACCTT 1300

45 TGAGGTCACT GCATTGAACG GGGTATCCTC CTTAGCCACG GGGCCCGTCC 1350

CATTTGAGCC TGTCAATGTC ACCACTGACC GAGAGGTACC TCCTGCAGTG 1400

50 TCTGACATCC GGGTGACGCG GTCCTCACCC AGCAGCTTGA GCCTGGCCTG 1450

55 GGCTGTTCCC CGGGCACCCA GTGGGGCTGT GCTGGACTAC GAGGTCAAAT 1500

ACCATGAGAA GGGCGCCGAG GGTCCCAGCA GCGTGCGGTT CCTGAAGACG 1550

60 TCAGAAAACC GGGCAGAGCT GCGGGGGCTG AAGCGGGGAG CCAGCTACCT 1600

GGTGCAAGTA CGGGCGCGCT CTGAGGCCGG CTACGGGCCC TTCGGCCAGG 1650

AACATCACAG CCAGACCCAA CTGGATGAGA GCGAGGGCTG GCGGGAGCAG 1700

5 CTGGCCCTGA TTGCGGGCAC GGCAGTCGTG GGTGTGGTCC TGGTCCTGGT 1750

GGTCATTGTG GTCGCAGTTC TCTGCCTCAG GAAGCAGAGC AATGGGAGAG 1800

10 AAGCAGAATA TTCGGACAAA CACGGACAGT ATCTCATCGG ACATGGTACT 1850

AAGGTCTACA TCGACCCCTT CACTTATGAA GACCCTAATG AGGCTGTGAG 1900

15 GGAATTTGCA AAAGAGATCG ATGTCTCCTA CGTCAAGATT GAAGAGGTGA 1950

20 TTGGTGCAGG TGAGTTTGGC GAGGTGTGCC GGGGGCGGCT CAAGGCCCCA 2000

GGGAAGAAGG AGAGCTGTGT GGCAATCAAG ACCCTGAAGG GTGGCTACAC 2050

25 GGAGCGGCAG CGGCGTGAGT TTCTGAGCGA GGCCTCCATC ATGGGCCAGT 2100

TCGAGCACCC CAATATCATC CGCCTGGAGG GCGTGGTCAC CAACAGCATG 2150

30 CCCGTCATGA TTCTCACAGA GTTCATGGAG AACGGCGCCC TGGACTCCTT 2200

35 CCTGCGGCTA AACGACGGAC AGTTCACAGT CATCCAGCTC GTGGGCATGC 2250

TGCGGGGCAT CGCCTCGGGC ATGCGGTACC TTGCCGAGAT GAGCTACGTC 2300

40 CACCGAGACC TGGCTGCTCG CAACATCCTA GTCAACAGCA ACCTCGTCTG 2350

CAAAGTGTCT GACTTTGGCC TTTCCCGATT CCTGGAGGAG AACTCTTCCG 2400

45 ATCCACCTA CACGAGCTCC CTGGGAGGAA AGATTCCCAT CCGATGGACT 2450

50 GCCCCGGAGG CCATTGCCTT CCGGAAGTTC ACTTCGCCA GTGATGCCTG 2500

GAGTTACGGG ATTGTGATGT GGGAGGTGAT GTCATTTGGG GAGAGGCCGT 2550

55 ACTGGGACAT GAGCAATCAG GACGTGATCA ATGCCATTGA ACAGGACTAC 2600

CGGCTGCCCC CGCCCCAGA CTGTCCCACC TCCCTCCACC AGCTCATGCT 2650

60 GGAATGCTTGG CAGAAAGACC GGAATGCCCC GCCCCGCTTC CCCCAGGTGG 2700



TCAGCGCCCT GGACAAGATG ATCCGGAACC CCGCCAGCCT CAAAATCGTG 2750

5 GCGCGGGAGA ATGGCGGGG CTCACACCCT CTCCTGGACC AGCGGCAGCC 2800

TCACTACTCA GCTTTTGGCT CTGTGGGCGA GTGGCTTCGG GCCATCAAAA 2850

10 TGGGAAGATA CGAAGAAAGT TTCGCAGCCG CTGGCTTTGG CTCCTTCGAG 2900

CTGGTCAGCC AGATCTCTGC TGAGGACCTG CTCCGAATCG GAGTCACTCT 2950

15 GCGCGGACAC CAGAAGAAAA TCTTGGCCAG TGTCCAGCAC ATGAAGTCCC 3000

AGGCCAAGCC GGGAAACCCG GGTGGGACAG GAGGACCGGC CCCGCAGTAC 3050

20 TGACCTGCAG GAACTCCCCA CCCAGGGAC ACCGCCTCCC CATTTTCCGG 3100

GGCAGAGTGG GGA CTCACAG AGGCCCCCAG CCCTGTGCCC CGCTGGATTG 3150

CACTTTGAGC CCGTGGGGTG AGGAGTTGGC AATTTGGAGA GACAGGATTT 3200

30 GGGGGTTCTG CCATAATAGG AGGGGAAAAT CACCCCCCAG CCACCTCGGG 3250

GAACTCCAGA CCAAGGGTGA GGGCGCCTTT CCCTCAGGAC TGGGTGTGAC 3300

35 CAGAGGAAAA GGAAGTGCCC AACATCTCCC AGCCTCCCCA GGTGCCCCC 3350

TCACCTTGAT GGGTGCGTTC CCGCAGACCA AAGAGAGTGT GACTCCCTTG 3400

CCAGCTCCAG AGTGGGGGGG CTGTCCCAGG GGGCAAGAAG GGGTGTGAGG 3450

45 GCCCAGTGAC AAAATCATTG GGGTTTGTAG TCCCAACTTG CTGCTGTCAC 3500

CACCAAATC AATCATTTTT TTCCCTTGTA AATGCCCTC CCCAGCTGC 3550

50 TGCCCTCATA TTGAAGGTTT TTGAGTTTGT TTTTGGTCT TAATTTTCT 3600

CCCCGTCCC TTTTGTTC TTCGTTTGT TTTTCTACCG TCCTTGTCAT 3650

AACTTTGTGT TGGAGGGAAC CTGTTTCACT ATGGCCTCCT TTGCCAAGT 3700

60 TGAAACAGGG GCCCATCATC ATGTCTGTTT CCAGAACAGT GCCTTGGTCA 3750

TCCCACATCC CCGGACCCG CCTGGGACCC CCAAGCTGTG TCCTATGAAG 3800

GGGTGTGGG TGAGGTAGTG AAAAGGGCGG TAGTTGGTGG TGAACCCAG 3850

5 AAACGGACGC CGGTGCTTGG AGGGGTCTT AAATTATATT TAAAAAGTA 3900

ACTTTTGTGTA TAAATAAAAG AAAATGGGAC GTGTCCCAGC TCCAGGGGTA 3950

10

AAAAAAAAA AAAAAAAAA 3969

15 (2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 987 amino acids

(B) TYPE: amino acid

20 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

25	Met	Glu	Leu	Arg	Val	Leu	Leu	Cys	Trp	Ala	Ser	Leu	Ala	Ala	Ala	1	5	10	15
	Leu	Glu	Glu	Thr	Leu	Leu	Asn	Thr	Lys	Leu	Glu	Thr	Ala	Asp	Leu	20	25	30	
30	Lys	Trp	Val	Thr	Phe	Pro	Gln	Val	Asp	Gly	Gln	Trp	Glu	Glu	Leu	35	40	45	
	Ser	Gly	Leu	Asp	Glu	Glu	Gln	His	Ser	Val	Arg	Thr	Tyr	Glu	Val	50	55	60	
35	Cys	Asp	Val	Gln	Arg	Ala	Pro	Gly	Gln	Ala	His	Trp	Leu	Arg	Thr	65	70	75	
40	Gly	Trp	Val	Pro	Arg	Arg	Gly	Ala	Val	His	Val	Tyr	Ala	Thr	Leu	80	85	90	
	Arg	Phe	Thr	Met	Leu	Glu	Cys	Leu	Ser	Leu	Pro	Arg	Ala	Gly	Arg	95	100	105	
45	Ser	Cys	Lys	Glu	Thr	Phe	Thr	Val	Phe	Tyr	Tyr	Glu	Ser	Asp	Ala	110	115	120	
	Asp	Thr	Ala	Thr	Ala	Leu	Thr	Pro	Ala	Trp	Met	Glu	Asn	Pro	Tyr	125	130	135	
50	Ile	Lys	Val	Asp	Thr	Val	Ala	Ala	Glu	His	Leu	Thr	Arg	Lys	Arg	140	145	150	
	Pro	Gly	Ala	Glu	Ala	Thr	Gly	Lys	Val	Asn	Val	Lys	Thr	Leu	Arg	155	160	165	
55	Leu	Gly	Pro	Leu	Ser	Lys	Ala	Gly	Phe	Tyr	Leu	Ala	Phe	Gln	Asp	170	175	180	
60	Gln	Gly	Ala	Cys	Met	Ala	Leu	Leu	Ser	Leu	His	Leu	Phe	Tyr	Lys	185	190	195	
	Lys	Cys	Ala	Gln	Leu	Thr	Val	Asn	Leu	Thr	Arg	Phe	Pro	Glu	Thr	200	205	210	

	Val	Pro	Arg	Glu	Leu	Val	Val	Pro	Val	Ala	Gly	Ser	Cys	Val	Val	
					215										220	
	Asp	Ala	Val	Pro	Ala	Pro	Gly	Pro	Ser	Pro	Ser	Leu	Tyr	Cys	Arg	
5					230										235	
	Glu	Asp	Gly	Gln	Trp	Ala	Glu	Gln	Pro	Val	Thr	Gly	Cys	Ser	Cys	
					245										250	
10	Ala	Pro	Gly	Phe	Glu	Ala	Ala	Glu	Gly	Asn	Thr	Lys	Cys	Arg	Ala	
					260										265	
	Cys	Ala	Gln	Gly	Thr	Phe	Lys	Pro	Leu	Ser	Gly	Glu	Gly	Ser	Cys	
					275										280	
15	Gln	Pro	Cys	Pro	Ala	Asn	Ser	His	Ser	Asn	Thr	Ile	Gly	Ser	Ala	
					290										295	
	Val	Cys	Gln	Cys	Arg	Val	Gly	Tyr	Phe	Arg	Ala	Arg	Thr	Asp	Pro	
20					305										310	
	Arg	Gly	Ala	Pro	Cys	Thr	Thr	Pro	Pro	Ser	Ala	Pro	Arg	Ser	Val	
					320										325	
25	Val	Ser	Arg	Leu	Asn	Gly	Ser	Ser	Leu	His	Leu	Glu	Trp	Ser	Ala	
					335										340	
	Pro	Leu	Glu	Ser	Gly	Gly	Arg	Glu	Asp	Leu	Thr	Tyr	Ala	Leu	Arg	
					350										355	
30	Cys	Arg	Glu	Cys	Arg	Pro	Gly	Gly	Ser	Cys	Ala	Pro	Cys	Gly	Gly	
					365										370	
	Asp	Leu	Thr	Phe	Asp	Pro	Gly	Pro	Arg	Asp	Leu	Val	Glu	Pro	Trp	
35					380										385	
	Val	Val	Val	Arg	Gly	Leu	Arg	Pro	Asp	Phe	Thr	Tyr	Thr	Phe	Glu	
					395										400	
40	Val	Thr	Ala	Leu	Asn	Gly	Val	Ser	Ser	Leu	Ala	Thr	Gly	Pro	Val	
					410										415	
	Pro	Phe	Glu	Pro	Val	Asn	Val	Thr	Thr	Asp	Arg	Glu	Val	Pro	Pro	
					425										430	
45	Ala	Val	Ser	Asp	Ile	Arg	Val	Thr	Arg	Ser	Ser	Pro	Ser	Ser	Leu	
					440										445	
	Ser	Leu	Ala	Trp	Ala	Val	Pro	Arg	Ala	Pro	Ser	Gly	Ala	Val	Leu	
50					455										460	
	Asp	Tyr	Glu	Val	Lys	Tyr	His	Glu	Lys	Gly	Ala	Glu	Gly	Pro	Ser	
					470										475	
55	Ser	Val	Arg	Phe	Leu	Lys	Thr	Ser	Glu	Asn	Arg	Ala	Glu	Leu	Arg	
					485										490	
	Gly	Leu	Lys	Arg	Gly	Ala	Ser	Tyr	Leu	Val	Gln	Val	Arg	Ala	Arg	
					500										505	
60	Ser	Glu	Ala	Gly	Tyr	Gly	Pro	Phe	Gly	Gln	Glu	His	His	Ser	Gln	
					515										520	
															525	

	Thr Gln Leu Asp	Glu Ser Glu Gly Trp	Arg Glu Gln Leu Ala	Leu
		530	535	540
5	Ile Ala Gly Thr	Ala Val Val Gly Val	Val Leu Val Leu Val	Val
		545	550	555
	Ile Val Val Ala	Val Leu Cys Leu Arg	Lys Gln Ser Asn Gly	Arg
		560	565	570
10	Glu Ala Glu Tyr	Ser Asp Lys His Gly	Gln Tyr Leu Ile Gly	His
		575	580	585
	Gly Thr Lys Val	Tyr Ile Asp Pro Phe	Thr Tyr Glu Asp Pro	Asn
		590	595	600
15	Glu Ala Val Arg	Glu Phe Ala Lys Glu	Ile Asp Val Ser Tyr	Val
		605	610	615
20	Lys Ile Glu Glu	Val Ile Gly Ala Gly	Glu Phe Gly Glu Val	Cys
		620	625	630
	Arg Gly Arg Leu	Lys Ala Pro Gly Lys	Lys Glu Ser Cys Val	Ala
		635	640	645
25	Ile Lys Thr Leu	Lys Gly Gly Tyr Thr	Glu Arg Gln Arg Arg	Glu
		650	655	660
	Phe Leu Ser Glu	Ala Ser Ile Met Gly	Gln Phe Glu His Pro	Asn
		665	670	675
30	Ile Ile Arg Leu	Glu Gly Val Val Thr	Asn Ser Met Pro Val	Met
		680	685	690
35	Ile Leu Thr Glu	Phe Met Glu Asn Gly	Ala Leu Asp Ser Phe	Leu
		695	700	705
	Arg Leu Asn Asp	Gly Gln Phe Thr Val	Ile Gln Leu Val Gly	Met
		710	715	720
40	Leu Arg Gly Ile	Ala Ser Gly Met Arg	Tyr Leu Ala Glu Met	Ser
		725	730	735
	Tyr Val His Arg	Asp Leu Ala Ala Arg	Asn Ile Leu Val Asn	Ser
		740	745	750
45	Asn Leu Val Cys	Lys Val Ser Asp Phe	Gly Leu Ser Arg Phe	Leu
		755	760	765
50	Glu Glu Asn Ser	Ser Asp Pro Thr Tyr	Thr Ser Ser Leu Gly	Gly
		770	775	780
	Lys Ile Pro Ile	Arg Trp Thr Ala Pro	Glu Ala Ile Ala Phe	Arg
		785	790	795
55	Lys Phe Thr Ser	Ala Ser Asp Ala Trp	Ser Tyr Gly Ile Val	Met
		800	805	810
	Trp Glu Val Met	Ser Phe Gly Glu Arg	Pro Tyr Trp Asp Met	Ser
		815	820	825
60	Asn Gln Asp Val	Ile Asn Ala Ile Glu	Gln Asp Tyr Arg Leu	Pro
		830	835	840

	Pro	Pro	Pro	Asp	Cys	Pro	Thr	Ser	Leu	His	Gln	Leu	Met	Leu	Asp
					845					850					855
5	Cys	Trp	Gln	Lys	Asp	Arg	Asn	Ala	Arg	Pro	Arg	Phe	Pro	Gln	Val
					860					865					870
	Val	Ser	Ala	Leu	Asp	Lys	Met	Ile	Arg	Asn	Pro	Ala	Ser	Leu	Lys
					875					880					885
10	Ile	Val	Ala	Arg	Glu	Asn	Gly	Gly	Ala	Ser	His	Pro	Leu	Leu	Asp
					890					895					900
	Gln	Arg	Gln	Pro	His	Tyr	Ser	Ala	Phe	Gly	Ser	Val	Gly	Glu	Trp
					905					910					915
15	Leu	Arg	Ala	Ile	Lys	Met	Gly	Arg	Tyr	Glu	Glu	Ser	Phe	Ala	Ala
					920					925					930
	Ala	Gly	Phe	Gly	Ser	Phe	Glu	Leu	Val	Ser	Gln	Ile	Ser	Ala	Glu
20					935					940					945
	Asp	Leu	Leu	Arg	Ile	Gly	Val	Thr	Leu	Ala	Gly	His	Gln	Lys	Lys
					950					955					960
25	Ile	Leu	Ala	Ser	Val	Gln	His	Met	Lys	Ser	Gln	Ala	Lys	Pro	Gly
					965					970					975
	Thr	Pro	Gly	Gly	Thr	Gly	Gly	Pro	Ala	Pro	Gln	Tyr			
					980					985		987			

30 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

40 Asp Lys Thr His Thr Cys Pro Pro Cys Pro

1 5 10

CLAIMS

1. Isolated hepatoma transmembrane kinase receptor ligand (Htk ligand).
- 5 2. The Htk ligand of claim 1, wherein the Htk ligand is antigenically active.
3. The Htk ligand of claim 1, wherein the Htk ligand is biologically active.
4. The Htk ligand of claim 1, comprising human Htk ligand.
- 10 5. The Htk ligand of claim 1, comprising a protein molecule including an amino acid sequence having at least 80% sequence identity with the amino acid sequence shown at positions 28 through 336, inclusive, of Figure 1A, and comprising biologically active Htk ligand.
6. The Htk ligand of claim 1, comprising a protein molecule  
15 including an amino acid sequence having at least 80% sequence identity with the amino acid sequence shown at positions 25 through 333, inclusive, of Figure 2, and comprising biologically active Htk ligand.
7. Soluble hepatoma transmembrane kinase receptor ligand (Htk ligand).
- 20 8. A composition comprising the Htk ligand of claim 1 and a pharmaceutically acceptable carrier.
9. A composition comprising the soluble Htk ligand of claim 7 and a pharmaceutically acceptable carrier.
10. An isolated nucleic acid molecule encoding the Htk ligand of  
25 claim 1.
11. The isolated nucleic acid molecule of claim 9 having a nucleic acid sequence selected from the group consisting of:
  - (a) residues 242 through 1168, inclusive, of the nucleic acid sequence shown in Figure 1A;
  - 30 (b) a sequence corresponding to the sequence of (a) within the scope of degeneracy of the genetic code; and
  - (c) a sequence which hybridizes with a sequence complementary to the sequence from (a) or (b) under stringent conditions and which codes for a biologically  
35 active Htk ligand.
12. The isolated nucleic acid molecule of claim 9 having a nucleic acid sequence selected from the group consisting of:
  - (a) residues 104 through 1030, inclusive, of the nucleic acid sequence shown in Figure 2;
  - 40 (b) a sequence corresponding to the sequence of (a) within the scope of degeneracy of the genetic code; and
  - (c) a sequence which hybridizes with a sequence complementary to the sequence from (a) or (b) under

stringent conditions and which codes for a biologically active Htk ligand.

13. An isolated nucleic acid molecule encoding the Htk ligand of claim 7.

5 14. A vector comprising the nucleic acid molecule of claim 10 operably linked to control sequences recognized by a host cell transformed with the vector.

15. A host cell comprising the vector of claim 14.

10 16. A method for preparing Htk ligand comprising culturing a host cell transfected to express the nucleic acid of claim 10 and recovering the Htk ligand from the host cell culture.

17. A chimeric polypeptide comprising a hepatoma transmembrane kinase receptor ligand (Htk ligand) amino acid sequence fused to an immunoglobulin sequence.

15 18. The chimeric polypeptide of claim 17 comprising a fusion of a Htk ligand extracellular domain sequence to an immunoglobulin constant domain sequence.

19. The chimeric polypeptide of claim 18 wherein said constant domain sequence is that of an immunoglobulin heavy chain.

20 20. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the chimeric polypeptide of claim 17.

21. A chimeric polypeptide comprising a hepatoma transmembrane kinase receptor ligand (Htk ligand) amino acid sequence fused to a tag polypeptide sequence.

25 22. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the chimeric polypeptide of claim 21.

23. A method for activating a tyrosine kinase domain of a hepatoma transmembrane kinase receptor (Htk receptor) comprising contacting an extracellular domain of the Htk receptor with the Htk ligand of claim 1.

30 24. A method for activating a tyrosine kinase domain of a hepatoma transmembrane kinase receptor (Htk receptor) comprising contacting an extracellular domain of the Htk receptor with the soluble Htk ligand of claim 7.

35 25. A monoclonal antibody which binds to the Htk ligand of claim 1.

```

1 CCCAGCGCTCCGCGCGCTGAGGGACGCGCAGGGTGAGCGCACCTGGCCTCGGCGACCGCGGGAGCGCGCGCGGTGTCCGCCCCGAGGATTGGGGGT

101 CGCTGCCCGCGCGCGGTCCCAACGCGTCCCGAGTGCGCAGAACTGGGAGCGGCTTGGGCATGGCCCATGGCCCCGGTCCAGGAGGGACTCTGTGTGGAAGT
      M A M A R S R R D S V W K Y
      10

201 ACTGTTGGGACTTTTGATGGTTTGTGCGAGAACTGCGATCTCCAGATCGATAGTTTGTAGAGCCTATCTACTGGAATTCCTCGAACTCCCAAATTTCTTACC
      C W G L L M V L C R T A I S R S I V L E P I Y W N S S N S K F L P
      20      30      40

301 CGGACAAGGCCTGGTACTATACCCACAGATAGGAGACAAATTTGGATATTATTGCCCCCAAGTGGACTCTAAAACCTGTGGCCAGTATGAATATATATAA
      G Q G L V L Y P Q I G D K L D I I C P K V D S K T V G Q Y E Y Y K
      50      60      70

401 GTTTATATGTTGATAAAGACCAAGCAGACAGATGCACAATTAAGAAGGAGATAACCCCGCTGTCTCAACTGTGCCAGACCAGACCAGATGTGAAATTCA
      V Y M V D K D Q A D R C T I K K E N T P L L N C A R P D Q D V K F T
      80      90      100      110

501 CCATCAAGTTTCAAGAATTCAGCCCTAACCTCTGGGGTCTAGAATTTCAGAAGAACAAGATTACTACATTATATCTACATCAAAATGGGTCTTTGGAGGG
      I K F Q E F S P N L W G L E F Q K N K D Y Y I I S T S N G S L E G
      120      130      140

601 CCTGGATNACCAGGAGGGGGTGTGCCAGACACAGAGCCATGAAGATCCTCATGAAAGTTGGACAAGATGCAAGTTCTGTGGATCAGCCAGGAATCAC
      L D N Q E G G V C Q T R A M K I L M K V G Q D A S S A G S A R N H
      150      160      170      180

701 GGTCACAAGACGTCCAGAGCTAGAAGCTGGTACAAATGGGAGAAGTTCAACAACAAGTCCCTTTGTGAAGCCAAATCCAGGTTCTAGCACCGATGGCA
      G P T R R P E L E A G T N G R S S T T S P F V K P N P G S S T D G N
      190      200      210

801 ACAGCGGGGGCATTCGGGGAACAATCTCCTGGGTTCCGAAGTGGCCTTATTTCGAGGGATCGCATCAGGATGATCATCTTTCATCGTCATCATCATCAC
      S A G H S G N N L L G S E V A L F A G I A S G C I I F I V I I I T
      220      230      240

```

FIG. 1A



2/11

250 260 270 280  
L V V L L L K Y R R R R K H S P Q H T T T L S L S T L A T P K R  
901 TTTGGTGGTGTGCTGCTCAAGTACCGCAGGAGACACCGCAAAACACTCTCCACAGCACACGACCACGCTGTCTCTCAGCACACTGGCCACGCCCCAAGCGA  
290 300 310  
G G N N N G S E P S D V I I P L R T A D S V F C P H Y E K V S G D Y  
1001 GGTGGCAACAATGGCTCGGAGCCCAAGTGACGTTATCATACCACTAAGGACTGCAGACAGCGTCTTCTGCCCCGCACTACGAGAAGGTCAGCGGGGACT  
320 330  
G H P V Y I V Q E M P P Q S P A N I Y Y K V O  
1101 ATGGGCACCCGGTGTACATCGTGCAGGAGATGCCCCCACAGAGTCTCTGCCAACATTTACTACAAGGTCAGAGCCTGAGCCTGAGACCTGCGCCTCCCAAGGGAAC  
1201 TCGCACCTTGTTCTTGGGCACGCAGGGACTGCCTGAGCCTGGCTGTGGGGGCAGGATGCCTCTGGAAGAGCCTGGATCTGGACAGTTTGTAGTCTGTA  
1301 GCTTTTCCGACCCCTGGGACCCACAGACCCCTCCCCCGGAAGCTGGAAGACTGCTAGGAGATCCCCACTTGGACTGCCCGCGGCCCCACGCGGACCTCCAAAGCC  
1401 ATGCACCCAGCCACTCAGGCCTCTGCAGAGCCCGGGGAGGACACGGTAGGCTATGGATGGCGCAGCAGCATCTTAGGAGAAGGTTGCGCACCCAGGCCCGGC  
1501 CCCTGCCTCCACGTTTCTGCGGTGCACACTGGACTTATCACTTGGACCTCGGGTTCAGTAAGGTTTTCAAAGATCTCTAGTGTCTTAGTCCCTCACTCACT  
1601 CACTCACTCACTCACTCTTCTTGTGCCAGGGCTCTGCAGCAAAACTCCCTAGACCCCTCACTCCACGTACTGCATCATTTACGGGACACTCACCCACAGAGT  
1701 CCCAGCTCCACCCCTTACACCAAGATCAAAATTAGATGGGTATTAGGTACAGAAGAACCCTGCTTGCCCTGGAGGCCCGGGTTCAGCCGGGAAGCGCAGATGTG  
1801 TGGAGGAGTGAGGAGGTGCTGGCTGCCACGGCAGGTCAAGGCTGCTTGCTGCCCTGGAGCATAGTAGGGATGCAGGAAGGAAATAGATAATGCTTTTGGT

FIG. 1B

3/11

1901 TTTTCTGAGAGGACAGAGACAGGTGGGAGGTGACTGCTGGTGGGAGCCCTTTCACTACACACAGCTATGCAGCAGGGAATCAAAAGTCCTCT  
2001 CTCCTGCGGGGAACAAAGGGGCCATTGTTGTGAAAGGACCAGCTAGAGCACAGAGGGAGAGGGCAGGCCCTCCGGTGAAAGTGCTGGGGCAGAACTGCAGAG  
2101 GTACTGGAAATAAAAAGCGCAGCGCAGAGCTGTGGGAGAGTCCGTCTGCTTTTGGGAGATGTTTTAAGCAGACTCAGCTGCTATATTACCACGTTTTTTATT  
2201 AAAAACACAGGGAAAGCATTTAGGAGAAAGCAGAGAGCCAAATCTGACCTAGAAAGTTGAAAAGCCAAAGGTCAAAACAGGCTGTAAAGTCCATCACCACTG  
2301 AGGTTATTGGAGAAATTCATATTAGGAAAGGCAGGTGAGATTCCTCCAGGCCCCCAATAAGTGTCCCCCTTCCCCCTGATTGAGCCCTTACACGTTGGTTTTT  
2401 GGTTTATGGCCGTGCTGTCCGGGCTCCAAAGGCAGTACCCGGGCTCCATGTCAAAAGCAAAAGCACACATGGCCCCACCTCTTAGAGTCCCTTGAGATGGAAGTA  
2501 AGTTATGCCCGGAAGGAAAGGCGAAGATAGGACATATTTATAATAGGTGTATAGAACAACAAGGGATATAAAATGAAAGATTTTTTACTAATATATATTTT  
2601 AAGATTACACACAATACACACCAGAAAGCGTGGAGTTCGGTGGTGGTGGTCCGTGGTGGTGGTGAATAAAGTGACCCCGCGCTTAGTGCTTTTAAAAA  
2701 GTGAAAGATTGGGTAGCTACTCCCCGAAACGTACCAATAGCAAGAAAAAGTATCCATAATGAGAGCAAAATGGCAAAAAATAACACGGTCCCTGCGGGAATCTC  
2801 GCAGAAACCGTAGACTAGGAATGCCAGCCCCCAATTTGATGTGACCTTGCCCCGGGTTAGACAATGATAAAATGCGCTGGCCCTTATTTTCTGTGTGGG  
2901 TTTTCCCTTGCCCTTATGGGCTGAAGTGTTCCTAGAAATTTAGCAGGTACACTGAGGGGATTCAGTTTAACGTGGGTCCCTCCTCTCCTACCCCCA  
3001 TCCCTGCCCTTCCAGAGAAATAACAGGAAGCCCTTCTTTTTTTTTTTTTTTTTTAAGTGCTATGCAAAAAGAGACATCTTTAACAGAGTCTCTGTACTATGGTAA

FIG. 1C

33101 CATTTTGCCTTCTCTGAATTGGGAGGAAATAAAAAATTGTAATGACAGCATTTTGAAGGTTCTCAGACCTCCAGTGAGTACCTGCAAAAAATGAGTTGTGCACAGA

33201 GATTATCCCTACTTCTCAAAACCTGAAAAATGATGTTGGTTCGATGTGTGTGTGTATGAGTGGGTGTGTGGTACATGTGTGTACATATATGTATATAAT

33301 ATATATCTCCAGTATATATTATATATATCTATATCATATTTCTGTGGAGGGTTGCCATGGCAATCAACTGCAGTACATATGTAGTTCTTTCCATCACCCCT

33401 AACCTCTCTGCGCATTCACACAAGAGTTTCTTGTAAGCCATCAAAAGTTAATTTCTAGGGGAGAGGGATGAGCGGGGAGACATGGGAAACCGTCTGTAT

33501 TTTAATGAAATCAAAATGCTGTGTGTCAATCGGTTGGCTACGTTTGGTTCTATGCTAAACTGTGAAGAATCGGATGAATTGATGAAGAGTTGAGTTACCTGC

33601 AACCCATTGAGAAAGTGCTCCTGTGCGTCTGTTTGTGTCTGGTGCAGAAAAATGACAAATCTACCAACTGTCCCCCTTATTTTGGAGTTGGTTTCAGCTTTTGGAAA

33701 GTTACTGTAAATGCCCTTGCTTGATTATCATCCCTAGTCACCTGACTTCGGAGCTTGACCCATCGTGTTTTAAAGTGAAGACGCTGTAAATAGGTTTCAGAT

33801 CTTACCGTCTATGGATTCCGGGTGTACAGTAGCCTTATTACCTTTTAAATAAAAAATACACATGAAAAACGAGACAGTAATGGCTTTTCTTACCCAGATTG

33901 TGTACATAGAGCAATGTTGGTTTTTTTATAAAGTCTAAAGCAAGATGTTTTGTATATAAAATCTGAATTTTGCATGTATTTAGCTACAGCTTTTAAACGGCAGT

34001 GTCATCCCCCTTGCACCTGTAATGAGGAAAAAAGGTAATAAAGGTTGCCAAATGCTGCATATTTGTGCGCGTAATTTATGTACCATGAATATTTTATTTT

34101 AATTTCGTTGTCCAAATTGTGAAGTAACACAGTATTATGCTTGAGTTATAAATATTTTTTCTTTGTTTATTTTAAATAGCCTGTTCATAGGTTTTTTTTTTTAA

34201 TCTGCTTTAGTTCCACATGACAGTTAAGCCCCAGAAATGAGATCCGAGCAGCCACATTCACCGTCTGTTCAAAAATGAATTTGTTCTTTAAAAAATAAA

34301 ATATTTTTTTTCTATGGAAAAAATAAAAAAAGGGCGGCGCCG

FIG.ID

**FIG. 1D**

5/11

1 CTCGAGCGCGGAGCTGGAGTGGCTTCGCCATGGCTGTGAGAAGGGAAGTCCGTGTGGAAGTACTGCTGGGGTGTTTGTGATGGTTTATGCAAACTGCG  
20  
M A V R R D S V W K Y C W G V L M V L C R T A  
30  
I S K S I V L E P I Y W N S S N S K F L P G Q G L V L Y P Q I G D K  
50  
101 ATTTCCAAATCGATAGTTTATAGAGCCCTATCTATTGGAATTCCTCGAACTCCAAATTTCTACCTGGACAAGGACTGGTACTATACCCACAGATAGGAGACA  
70  
L D I I C P K V D S K T V G Q Y E Y Y K V Y M V D K D Q A D R C T  
90  
201 AATTGGATATTATTGCCCCCAAGTGGACTCTAAAACTGTGGCCAGTATGAATATTATAAGTTTATATGGTTGATAAAGACCAAGCAGACAGATGCAC  
110  
I K K E N T P L L N C A K P D Q D I K F T I K F Q E F S P N L W G  
120  
301 TATTAAGAAGGAAATACCCCTCTCCTCAACTGTGCCAAACCAGACCAAGATATCAATTCACCATCAAGTTTCAAGAATTCAGCCCTAACCTCTGGGGT  
140  
L E F Q K N K D Y Y I I S T S N G S L E G L D N Q E G V C Q T R A  
150  
401 CTAGAATTTCAGAAGAACAAGATTATTACATTATATCTACATCAAAATGGGTCTTTGGAGGGCCTGGATAACCAAGGAGGGGGTGTGCCAGACAAAG  
170  
M K I L M K V G Q D A S S A G S T R N K D P T R R P E L E A G T N  
180  
501 CCATGAAGATCCTCATGAAGTTGGACAAGATGCAAGTTCTGTGGATCAACCAGGAATAAAGATCCAAACAAGACGTCCAGAAGTCTAGAAAGCTGGTACAAA  
200  
G R S S T T S P F V K P N P G S S T D G N S A G H S G N N I L G S  
210  
601 TGGAGAAGTTCGACAACAAGTCCCTTTGTAAACCAATCCAGGTTCTAGCACAGACGGCAACAGCCCGGACATTCGGGGGAACAACATCCTCGGTTCC  
220  
230  
E V A L F A G I A S G C I I F I V I I I T L V V L L L K Y R R R H R  
240  
701 GAAGTGGCCTTATTGACGGGATGCTTCAGGATGCATCATCTTCATCGTTCATCATCAGCGTGGTGGTCTCTTGTGTAAGTACCGGAGGAGACACA  
250  
260  
K H S P Q H T T T L S L S T L A T P K R S G N N N G S E P S D I I  
270  
801 GGAAGCACTGCGCGCAGCACACGACCACGCTGTGGCTCAGCACACTGGCCACACCCCAAGCGCAGCGGCAACAACACCGGCTCAGAGCCCCAGTGACATTAT  
280  
290

FIG.2A

300 310 320  
I P L R T A D S V F C P H Y E K V S G D Y G H P V Y I V Q E M P P  
901 CATCCGCTAAGGACTGCGGACAGCGTCTTCTGCTCCTCACTACGAGAAGGTACGGGGACTACGGGCACCCGGTGATACATCGTCCAGGAGATGCCCCCG  
330  
Q S P A N I Y Y K V O  
1001 CAGAGCCCGCGAACATTTACTACAAGGTCTGAGAGGGACCTGGTGGTACCTGTGCTTTCCAGAGGACACCTAATGTCCCGATGCCCTTGAGGGT  
1101 TTGAGAGCCCGGTGCTGGAGAAATTGACTGAAGCACAGCACCGGGGAGAGGGACACTCCTCCTCGGAAGAGCCCGTGGCTGGACAGCTTACCTAGTGC  
1201 TTGTAGCATTCGGCCTTGGTGAACACACACGCTCCCTGGAAAGCTGGAAAGACTGTGCAGAAAGACGCCCATTCGGACTGCTGTGCCGCTGCCACGTCCTCT  
1301 CCTCGAAGCCATGTGCTGCGGTCACTCAGGCCCTCTGCAGAACCCAAAGGAAGACAGTGGTTTGTGGACGAGAGGGCTGTGAGCATCCTGGCAGGTGCCCC  
1401 AGGATGCCACGCCCTGGAAGGGCCGGCTTCTGCCTGGGGTGCAATTTCCCCCGCAGTGCAATACCGGACTTGTGTCAACGACCTCGGGCTAGTTAAGGTGTGC  
1501 AAAGATCTCTAGAGTTTAGTCCTTACTGTCTCACTCGTTCTGTACCCAGGGCTCTGCAGCACCTCACCTGAGACCTCCACATCTGCATCACTCA  
1601 TGGAACTCATGTCTGGAGTCCCCCTCCTCCAGCCGCTGSGCAACAACAGCTTCAGTCCATGGGTAATCCCGTTATAGAAATTGTGTTTGCTAACAAAGGTG  
1701 CCCTTTAGCCAGATGCTAGGCTGTCTGCGAAGAAGGCTAGGAGTTCTAGAAGGGAGTGGGGCTGGGAAAGGGCTGGCTGCAATTGCAGCTCACTGCTG  
1801 CTGCCCTCTGAAACAGAAAGTTGGAAAGGAAAAAAGCAATTAGGTAGCACAGCACTTTGGTTTTGCTGAGATCGAAAGAGGCCAGTAGGAGACAC  
1901 GACAGCACACAGTGGATTCCAGTGCATGGGGAGGCGGTCCAGCAGCTCGAG

FIG.2B

[illegible]

**FIG. 3**

1 TCGGCGTCCACCCCGGAGAGTACACCTGGGGGGGGGAGGGCCCCCAACTCACTTCGGATCCTACCCGAGTGAGGGGGCCCATGGAGCTCCG M E L R  
10 V L L C W A S L A A A L E E T L L L N T K L E T A D L K W V T F P Q 30  
101 GGTGCTGCTGCTGGCTTGGTGGCGCAGCTTTGGAAGAGACCCCTGCTGAACACAAAATTGGAACCTGCTGATCTGAAGTGGTGACATTCCTCCTCAG 40  
V D G Q W E E L S G L D E E Q H S V R T Y E V C D V Q R A P G Q A H 60  
201 GTGCAGGGCAGTGGGAGAACTGAGCGGCCTGGATGAGGAACAGACACAGCGTGGCAGCTACGAAATGTGTGACGTGCAGCGTGCCCCGGCCAGGCCCC 80  
W L R T G W V P R R G A V H V Y A T L R F T M L E C L S L P R A G 100  
301 ACTGGCTTCGCACAGTGGGTCCACGGGGCGCGCTCCACGTGTACGGCTTACGCCAGCTGGCTTCAACATGCTCGAGTGCCTGTCCCTGCTCGGGCTGG 120  
R S C K E T F T V F Y Y E S D A T A T A L T P A W M E N P Y I K 130  
401 GCGCTCTGCAAGGAGACCTTACCGTCTTCTACTATGAGAGCGATGGGACACGGCCACGGCCCTCAGCCAGCTGGATGGAGAACCCCTACATCAAG 140  
V D T V A A E H L T R K R P G A E A T G K V N V K T L R L G P L S K 160  
501 GTGGACACGCTGGCCCGGAGCATCTACCCGGAAGCGCCCTGGGGCGAGGCCACCGGAAGTGAATGTCAAGACGCTGCGTCTGGGACCGCTCAGCA 180  
A G F Y L A F Q D Q G A C M A L L S L H L F Y K K C A Q L T V N L 190  
601 AGGTGGCTTCTACCTGGCTTCCAGGACACAGGTGCTGCTGATGGCCCTGCTATCCCTGCACCTCTTCTACAAAAGTGGCCCCAGCTGACTGTGAACCT 210  
T R F P E T V P R E L V V P V A G S C V V D A V P A P G P S P S L 220  
701 GACTCGATTCCCGGAGACTGTGCTCGGAGTGTGTGTCCTGGCGGTAGTGGCTGTGGATGCGCTGCCCGCCCTGGCCCCCAGCCCGCTC 240  
Y C R E D G Q W A E Q P V T G C S C A P G F E A A E G N T K C R A C 250  
801 TACTGCGGTGAGGATGCCAGTGGGCGGAACAGCGGTACAGGGTGCAGTGTGCTCGGGGTTCGAGGCAGCTGAGGGGAACACCAAGTCCCGAGCCT 260  
A Q G T F K P L S G E G S C Q P C P A N S H S N T I G S A V C Q 270  
901 GTGCCAGGGCACCTTCAAGCCCTGTTCAGGAGAAGGTCTCTGCCAGCCATGCCCGCCCAATAGCCACTTAACACCATTGGATCAGCCGTCTGCCAGTG 280  
R V G Y F R A R T D P R G A P C T T P P S A P R S V V S R L N G S 290  
11001 CCGCGTGGGTACTTCCGGGCGCACAGACCCCGGGGTGCACCCCTGCACCACCCCTCCTTCGGCTCGGGGAGCGTGTGTTCCCGCCTGAACGGCTCC 310  
S L H L E W S A P L E S G G R E D L T Y A L R C R E C R P G G S 320  
1101 TCCCTGCACCTGGAATGGAGTGCCCCCTGGAGTCTGTGGCGGAGAGGACCTACCTACGCCCTCGCTGCGGGAGTGCCACCCCGGAGGCTCTCTGTG 330  
340 350 360 370

**FIG. 4A**

[illegible]

**FIG. 4B**



**FIG. 4C**

11/11




FIG. 5A




FIG. 5B